



Ambient Water Quality Criteria Recommendations

**Information Supporting the Development
of State and Tribal Nutrient Criteria**

Lakes and Reservoirs in Nutrient Ecoregion V



AMBIENT WATER QUALITY CRITERIA RECOMMENDATIONS

**INFORMATION SUPPORTING THE DEVELOPMENT OF STATE AND TRIBAL
NUTRIENT CRITERIA**

FOR

LAKES AND RESERVOIRS IN NUTRIENT ECOREGION V

South Central Cultivated Great Plains

including all parts of the States of:

*Montana, North Dakota, South Dakota, Nebraska, Wyoming, Colorado,
Kansas, Oklahoma, Texas, and New Mexico,*

and the authorized Tribes within the Ecoregion

U.S. ENVIRONMENTAL PROTECTION AGENCY

**OFFICE OF WATER
OFFICE OF SCIENCE AND TECHNOLOGY
HEALTH AND ECOLOGICAL CRITERIA DIVISION
WASHINGTON, DC**

DECEMBER 2001

FOREWORD

This document presents EPA's nutrient criteria for **Lakes and Reservoirs in Nutrient Ecoregion V**. These criteria provide EPA's recommendations to States and authorized Tribes for use in establishing their water quality standards consistent with section 303(c) of the Clean Water Act (CWA). Under section 303(c) of the CWA, States and authorized Tribes have the primary responsibility for adopting water quality standards as part of State or Tribal law or regulation. Federal regulations require State and Tribal standards to contain scientifically defensible water quality criteria that are protective of designated uses. EPA's recommended section 304(a) criteria are not laws or regulations; they are guidance that States and Tribes may use as a starting point in creating their own water quality standards.

The term "water quality criteria" is used in two sections of the CWA, section 304(a)(1) and section 303(c)(2). The term has a different impact in each section. On the one hand, in section 304, the term represents a scientific assessment of ecological and human health effects that EPA recommends to States and authorized Tribes for establishing water quality standards that ultimately provide a basis for controlling discharges or releases of pollutants or related parameters. On the other hand, in section 303, ambient water quality criteria are developed by States and Tribes as part of their water quality standards, to define the level of a pollutant (or in the case of nutrients, a condition) necessary to protect designated uses in ambient waters.

Quantified water quality criteria contained within State or Tribal water quality standards are essential to a water quality-based approach to pollution control. Whether expressed numerically or as quantified translations of narrative criteria within State or Tribal water quality standards, quantified criteria are critical for assessing attainment of designated uses and measuring progress toward meeting CWA goals.

EPA is developing section 304(a) water quality criteria for nutrients because States and Tribes consistently identify excessive levels of nutrients as a major reason that as many as half of the Nation's surface waters surveyed do not meet water quality objectives, such as full support of aquatic life. EPA expects to develop nutrient criteria that cover four major types of waterbodies—lakes and reservoirs, rivers and streams, estuarine and coastal areas, and wetlands—across 14 major ecoregions of the United States. EPA's section 304(a) criteria are intended to provide for the protection and propagation of aquatic life and recreation. To support the development of nutrient criteria, EPA has published and will continue to publish technical guidance manuals that describe a process for assessing nutrient conditions in the four waterbody types listed above.

EPA's section 304(a) water quality criteria for nutrients provide numeric water quality criteria and procedures to help establish quantified criteria within State or Tribal water quality standards. In the case of nutrients, EPA section 304(a) criteria establish values for causal variables (e.g., total nitrogen and total phosphorus) and response variables (e.g., Secchi depth and chlorophyll *a*). EPA believes that State and Tribal water quality standards need to include quantified endpoints for causal and response variables to provide sufficient protection of uses and to maintain downstream uses. These endpoints will most often be expressed as numeric water quality criteria or as procedures to translate a State or Tribal narrative criterion into a quantified endpoint.

States and authorized Tribes have several options in adopting these criteria. EPA recommends the following approaches, in order of preference:

1. Wherever possible, develop nutrient criteria that fully reflect local conditions and protect specific designated uses through the process described in EPA's technical guidance manuals for nutrient criteria development. Such criteria may be expressed either as numeric criteria or as procedures to translate a State or Tribal narrative criterion into a quantified endpoint in State or Tribal water quality standards.
2. Adopt EPA's section 304(a) water quality criteria for nutrients, either as numeric criteria or as procedures to translate a State or Tribal narrative nutrient criterion into a quantified endpoint.
3. Develop nutrient criteria protective of designated uses using other scientifically defensible methods and appropriate water quality data.

EPA developed the nutrient criteria recommendations in this document with the intent that they serve as a starting point for States and Tribes to develop more refined criteria, as appropriate, to reflect local conditions. The values presented in this document generally represent nutrient levels that protect against the adverse effects of nutrient overenrichment. They are based on the information that was available to the Agency at the time of this publication. EPA expects States and Tribes may have additional information and data that may be utilized in the refinement of these criteria. EPA offers to work with States and authorized Tribes to establish the necessary quantitative endpoints to reduce the excess nutrient inputs into our nation's waters and to prevent any further impairments.

Geoffrey H. Grubbs, Director
Office of Science and Technology

DISCLAIMER

This document provides technical guidance and recommendations to States, authorized Tribes, and other authorized jurisdictions to develop water quality criteria and water quality standards under the Clean Water Act (CWA) to protect against the adverse effects of nutrient overenrichment. Under the CWA, States and authorized Tribes are to establish water quality criteria to protect designated uses. State and Tribal decisionmakers retain the discretion to adopt approaches on a case-by-case basis that differ from this guidance when appropriate and scientifically defensible. Even though this document contains EPA's scientific recommendations regarding ambient concentrations of nutrients that will protect aquatic resource quality, it does not substitute for the CWA or EPA regulations, nor is it a regulation itself. Thus it cannot impose legally binding requirements on EPA, States, authorized Tribes, or the regulated community, and it might not apply to a particular situation or circumstance. EPA may change this guidance in the future.

EXECUTIVE SUMMARY

Nutrient Program Goals

EPA developed the National Strategy for the Development of Regional Nutrient Criteria (National Strategy) in June 1998. The strategy presents EPA's intentions to develop technical guidance manuals for four types of waters (lakes and reservoirs, rivers and streams, estuaries and coastal waters, and wetlands) and produce section 304(a) criteria for specific nutrient Ecoregions by the end of 2000. In addition, the Agency formed Regional Technical Assistance Groups (RTAGs), which include State and Tribal representatives working to develop more refined and localized nutrient criteria based on approaches described in the waterbody guidance manuals. This document presents EPA's current recommended criteria for total phosphorus (TP), total nitrogen (TN), chlorophyll *a*, and Secchi for lakes and reservoirs in Nutrient Ecoregion V, which were derived using the procedures described in the *Lakes and Reservoirs Nutrient Criteria Technical Guidance Manual* (U.S. EPA, 2000a).

EPA's ecoregional nutrient criteria address cultural eutrophication—the adverse effects of excess human-caused nutrient inputs. The criteria are empirically derived to represent surface waters that are minimally impacted by human activities and protective of aquatic life and recreational uses. The information contained in this document represents starting points for States and Tribes to develop (with assistance from EPA) more refined nutrient criteria.

In developing these criteria recommendations, EPA followed a process that included, to the extent they were readily available, the following critical elements:

- **Historical and recent nutrient data in Nutrient Ecoregion V.** Data sets from Legacy STORET, EPA Region 8 - Montana and Wyoming, EPA Region 8 - South Dakota, and EPA Region 8 - North Dakota were used to assess nutrient conditions from 1990 to 2000.
- **Reference sites/reference conditions in Nutrient Ecoregion V.** Reference conditions presented are based on 25th percentiles of all nutrient data, including a comparison of reference conditions for the Aggregate Ecoregion versus the subecoregions. States and Tribes are urged to determine their own reference sites for lakes and reservoirs at different geographic scales and to compare them to EPA's reference conditions.
- **Models employed for prediction or validation.** EPA did not identify any specific models to develop nutrient criteria. States and Tribes are encouraged to identify and apply appropriate models to support nutrient criteria development.
- **RTAG expert review and consensus.** EPA recommends that when States and Tribes prepare their nutrient criteria, they obtain the expert review and consent of the RTAG.
- **Downstream effects of criteria.** EPA encourages the RTAG to assess the potential effects of the proposed criteria on downstream water quality and uses.

In addition, EPA followed specific **QA/QC procedures** during data collection and analysis. All data were reviewed for duplications. All data were from ambient waters that were not located directly outside a permitted discharger. The following states indicated that their data were sampled and analyzed using either standard methods or EPA-approved methods: North Dakota, South Dakota, Montana, Wyoming, Colorado, Kansas, Oklahoma, and Texas. Nebraska indicated that standard or EPA-approved methods were used for some specific nutrient parameters.

The following tables contain a summary of aggregate and level III Ecoregion values for TN, TP, water column chlorophyll *a*, and Secchi.

BASED ON 25th PERCENTILES ONLY

Nutrient Parameters	Aggregate Nutrient Ecoregion V Reference Conditions
Total phosphorus (µg/L)	33
Total nitrogen (mg/L) (calculated)	0.56
Chlorophyll <i>a</i> (µg/L) (spectrophotometric method)	2.3
Secchi (m)	1.3

For subcoregions 25, 27, 32, and 42 the ranges of nutrient parameter reference conditions are as follows:

BASED ON 25th PERCENTILE ONLY

Nutrient Parameters	Range of Level III Subcoregions Reference Conditions
Total phosphorus (µg/L)	24-40
Total nitrogen (mg/L) (calculated)	0.50-1.36
Chlorophyll <i>a</i> (µg/L) (spectrophotometric method)	0.2-5.8
Secchi (m)	0.8-2.4

NOTICE OF DOCUMENT AVAILABILITY

This document is available electronically to the public through the Internet at <http://www.epa.gov/OST/standards/nutrient.html>. Requests for hard copies of the document should be made to EPA's National Service Center for Environmental Publications (NSCEP), 11029 Kenwood Road, Cincinnati, OH 45242; telephone (513) 489-8190 or toll free (800) 490-9198. Please refer to EPA document number EPA-822-B-01-010.

ACKNOWLEDGMENTS

The authors thankfully acknowledge the contributions of the following State and Federal reviewers: EPA Regions 6, 7, and 8; the States of North Dakota, South Dakota, Montana, Wyoming, Nebraska, Colorado, Kansas, Oklahoma, New Mexico, and Texas; the Tribes within the Ecoregion; EPA headquarters personnel from the Office of Wetlands, Oceans, and Watersheds, Office of Wastewater Management, Office of General Counsel, Office of Research and Development, and Office of Science and Technology. EPA also acknowledges the external peer review efforts of Paul Garrison, Wisconsin Department of Natural Resources; John Reuter, University of California, Davis; and Eugene Welch, University of Washington.

TABLE OF CONTENTS

Foreword	iii
Disclaimer	v
Executive Summary	vii
Notice of Document Availability	ix
Acknowledgments	x
List of Tables and Figures	xii
 1.0 Introduction	 1
 2.0 Best Use of This Information	 6
 3.0 Area Covered by This Document	 8
3.1 Description of Aggregate Ecoregion V	8
3.2 Geographical Boundaries of Aggregate Ecoregion V	9
3.3 Level III Subcoregions Within Aggregate Ecoregion V	11
3.4 Suggested Ecoregional Subdivisions or Adjustments	13
 4.0 Data Review for Lakes and Reservoirs in Aggregate Ecoregion V	 13
4.1 Data Sources	13
4.2 Historical Data from Aggregate Ecoregion V (TP, TN, Chl <i>a</i> , and Secchi)	13
4.3 QA/QC of Data Sources	13
4.4 Data for All Lakes and Reservoirs Within Aggregate Ecoregion V	14
4.5 Statistical Analysis of Data	14
4.6 Classification of Lake/Reservoir Type	21
4.7 Summary of Data Reduction Methods	22
 5.0 Reference Sites and Conditions in Aggregate Ecoregion V	 25
 6.0 Models Used to Predict or Verify Response Parameters	 25
 7.0 Framework for Refining Recommended Nutrient Criteria for Lakes and Reservoirs in Aggregate Ecoregion V	 26
7.1 Example Worksheet for Developing Aggregate Ecoregion and Subcoregion Nutrient Criteria	26
7.2 Setting Seasonal Criteria	27
7.3 When Data/Reference Conditions Are Lacking	27
7.4 Site-Specific Criteria Development	28
 8.0 Literature Cited	 28
 9.0 Appendices	 29
Appendix A. Descriptive Statistics Data Tables for Aggregate Ecoregion	A-1
Appendix B. Descriptive Statistics Data Tables for level III Subcoregions Within Aggregate Ecoregion	B-1
Appendix C. Quality Control/Quality Assurance Rules	C-1

LIST OF TABLES AND FIGURES

Tables

Table 1	Lake records for Aggregate Ecoregion V— South Central Cultivated Great Plains	16
Table 2	Reference conditions for Aggregate Ecoregion V lakes and reservoirs	17
Table 3a-d	Reference conditions for Ecoregion V lakes and reservoirs	18
Table 4	Changes in temperate lake attributes according to trophic state	21

Figures

Figure 1a	Fourteen nutrient ecoregions as delineated by Omernik (2000)	4
Figure 1b	Level III Ecoregions of the United States	5
Figure 2	Aggregate Ecoregion V	10
Figure 3	Aggregate Ecoregion V with level III Ecoregions shown	12
Figure 4	Sampling locations within each level III Ecoregion	15
Figure 5a	Illustration of data reduction process for lake data	23
Figure 5b	Illustration of reference condition calculation	24

1.0 INTRODUCTION

Background

Nutrients are essential to the health and diversity of surface waters. However, in excessive amounts nutrients cause eutrophication or hypereutrophication, which results in overgrowth of plant life and decline of the biological community. Excessive nutrients can also result in human health risks, such as the growth of harmful algal blooms, most recently manifested in the *Pfiesteria* outbreaks on the Gulf and East Coasts. Chronic nutrient overenrichment of a waterbody can lead to the following consequences: algal blooms, low dissolved oxygen, fish kills, overabundance of macrophytes, likely increased sedimentation and species shifts of both flora and fauna.

Historically, National Water Quality Inventories have repeatedly shown that nutrients are a major cause of ambient water quality use impairments. EPA's 1996 National Water Quality Inventory report identifies excessive nutrients as the leading cause of impairment in lakes and the second leading cause of impairment in rivers (behind siltation). In addition, nutrients were the second leading cause of impairments after siltation reported by the States in their 1998 lists of impaired waters. Where use impairment is documented, nutrients contribute roughly 25%-50% of the impairment nationally. The Clean Water Act (CWA) establishes that, wherever possible, water quality must provide for the protection and propagation of fish, shellfish, and wildlife, and recreation in and on the water and/or protecting the physical, chemical, and biological integrity of those waters. In adopting water quality standards, States and Tribes designate uses for their waters in consideration of these CWA goals, and establish water quality criteria that contain sufficient parameters to protect that integrity and those uses. To date, EPA has not published information and recommendations under section 304(a) for nutrients to assist States and Tribes in establishing numeric nutrient criteria to protect uses when adopting water quality standards.

In 1995, EPA gathered a set of national experts and asked them how best to deal with the national nutrient problem. The experts recommended that the Agency not develop single criteria values for phosphorus (P) or nitrogen (N) applicable to all waterbodies and regions of the country. Rather, they recommended that EPA put a premium on regionalization, develop guidance (assessment tools and control measures) for specific waterbodies and ecological regions across the country, and use reference conditions (conditions that reflect pristine or minimally impacted waters) as a basis for developing nutrient criteria.

With these suggestions as starting points, EPA developed the National Strategy for the Development of Regional Nutrient Criteria (National Strategy), published in June 1998. This strategy presented EPA's intentions to develop technical guidance manuals for four types of waters (lakes and reservoirs, rivers and streams, estuaries and coastal waters, and wetlands), and thereafter to publish section 304(a) criteria recommendations for specific nutrient Ecoregions. Technical guidance manuals for lakes/reservoirs and rivers/streams were published in April 2000 and July 2000, respectively. The technical guidance manual for estuaries/coastal waters was published in fall 2001, and the draft wetlands technical guidance manual will be published by December 2001. Each manual presents EPA's recommended approach for developing nutrient criteria values for a specific waterbody type. In addition, EPA is committed to working with

States and Tribes to develop more refined and localized nutrient criteria based on approaches described in the waterbody guidance manuals and this document.

Overview of the Nutrient Criteria Development Process

For each nutrient Ecoregion, EPA developed a set of recommendations for two causal variables (total nitrogen and total phosphorus) and two early indicator response variables (chlorophyll *a* [chl *a*] and Secchi). Other indicators such as dissolved oxygen, macrophyte or benthic algal growth or speciation, and other fauna and flora changes are also useful. However, the first four variables are considered to be the best suited for protecting designated uses.

The technical guidance manuals describe a process for developing nutrient criteria that involves consideration of five factors. The first of these is the Regional Technical Assistance Group (RTAG), which is a body of qualified regional specialists able to objectively evaluate all of the available evidence and select the value(s) appropriate to nutrient control in the water bodies of concern. These specialists may come from such disciplines as limnology, biology, or natural resources management—especially water resource management, chemistry, and ecology. The RTAG evaluates and recommends appropriate classification techniques, usually physical, for criteria determination within an ecoregional construct.

The second factor is the historical information available to establish a perspective of the resource base. This is usually data and anecdotal information available within the past 10-25 years. This information gives evidence about the background and enrichment trend of the resource.

The third factor is the existing reference condition, a selection of reference sites chosen to represent the least culturally impacted waters of the class at the present time. The data from these sites are combined and a value is selected to represent the reference condition, the best attainable, most natural condition of the resource base at this time.

The RTAG comprehensively evaluates these three elements to propose a candidate criterion (initially one each for TP, TN, chl *a*, and Secchi).

A fourth factor often employed is mechanistic or empirical models of the historical and reference condition data to better understand the condition of the resource.

The final element of the process is assessment by the RTAG of the likely downstream effects of the criterion. Will there be a negative, positive, or neutral effect on the downstream waterbody? If the RTAG judges that a negative effect is likely, then the proposed State/Tribal water quality criteria should be revised to ameliorate the potential for any adverse downstream effects.

Although States and authorized Tribes do not necessarily need to incorporate all five elements into their water quality criteria setting process (e.g., modeling may be significant in only some instances), the best assurance of a representative and effective criterion is a balanced incorporation of all five elements.

Because some parts of the country have naturally different soil and parent material nutrient content and different precipitation regimes, the application of the criterion development process should reflect this regional variation. Therefore, an ecoregional approach was chosen. Initially, the continental United States was divided into 14 separate Ecoregions of similar geographical characteristics and similar nutrient condition (Figure 1a). Ecoregions are defined as regions of relative homogeneity in ecological systems; they depict areas within which the mosaic of ecosystem components (biotic and abiotic as well as terrestrial and aquatic) is different from adjacent areas in a holistic sense. Geographic characteristics such as soils, vegetation, climate, geology, and land cover, are relatively similar within each Ecoregion (Omernik, 2000).

The nutrient Ecoregions are aggregates of EPA's hierarchical level III Ecoregions (see Figure 1b for a map of level III Ecoregions). As such, they are more generalized and less defined than level III Ecoregions. EPA determined that setting ecoregional criteria for the large-scale aggregates is not without its drawbacks: variability is high because of the lumping of many waterbody classes, seasons, and years worth of multipurpose data over a large geographic area. For these reasons, the Agency recommends that States and Tribes develop nutrient criteria at the level III ecoregional scale and at the waterbody-class scale, where those data are readily available. Data analyses and recommendations on both the large Aggregate Ecoregion scale and the more refined scales (level III Ecoregions and waterbody classes), where data were available to make such assessments, are presented for comparison and completeness of analysis.

Comparison of Nutrient Criteria to Biological Criteria

Biological criteria are quantitative expressions of the desired condition of the aquatic community. Such criteria can be based on data from sites that represent the least impacted attainable condition for a particular waterbody type in an Ecoregion, subecoregion, or watershed. EPA's nutrient criteria recommendations and biological criteria recommendations have many similarities in their basic approaches to development and data requirements. Both are empirically derived from statistical analysis of field-collected data and expert evaluation of current reference conditions and historical information. Both use direct measurements from the environment to integrate the effects of complex processes that vary according to type and location of waterbody. The resulting criteria recommendations, in both cases, are efficient uses of existing resources and are holistic indicators of the water quality necessary to protect uses.

States and authorized Tribes can develop and apply nutrient and biological criteria in tandem, with each providing important and useful information to interpret both the nutrient enrichment levels and the biological condition of sampled waterbodies. For example, using the same reference sites for both types of criteria can lead to efficiencies in both sample design and data analysis. In one effort, environmental managers can obtain information to support assessment of biological and nutrient condition, either through evaluating existing data sets or through designing and conducting a common sampling program. The traditional biological criteria variables of benthic invertebrate and fish sampling can be readily incorporated in a




Figure 1a. Fourteen nutrient Ecoregions as delineated by Omernik (2000). Ecoregions were based on geology, land use, ecosystem type, and nutrient conditions.




Figure 1b. Level III Ecoregions of the United States.

nutrient assessment. To investigate the effectiveness of this tandem approach, EPA has initiated pilot projects in both freshwater and marine environments to pursue the relationship between nutrient overenrichment and apparent declines in diversity of benthic invertebrates and fish.

2.0 BEST USE OF THIS INFORMATION

EPA recommendations published under section 304(a) of the CWA serve several purposes, including providing guidance to States and Tribes in adopting water quality standards for nutrients and ultimately controlling discharges or releases of pollutants. The recommendations also provide guidance to EPA when it determines that it is necessary to promulgate Federal water quality standards under section 303(c). Other uses include identification of overenrichment problems, management planning, project evaluation, and determination of status and trends of water resources.

State water quality inventories and listings of impaired waters consistently rank nutrient overenrichment as a top contributor to use impairments. EPA's water quality standards regulations at 40 CFR §131.11(a) require States and Tribes to adopt criteria that contain sufficient parameters and constituents to protect the designated uses of their waters. In addition, States and Tribes need quantifiable targets for nutrients to assess attainment of uses, develop water quality-based permit limits and source control plans, and establish targets for total maximum daily loads (TMDLs).

EPA expects States and Tribes to address nutrient overenrichment in their water quality standards and to build on existing State and Tribal efforts where possible. States and Tribes can address nutrient overenrichment through establishment of numerical criteria or use of narrative criteria statements (e.g., "free from excess nutrients that cause or contribute to undesirable or nuisance aquatic life or produce adverse physiological response in humans, animals, or plants"). In the case of narrative criteria, EPA expects that States and Tribes will establish procedures to quantitatively translate these statements for both assessment and source control purposes.

Ecoregional nutrient criteria are developed to represent surface waters that are minimally impacted by human activities and thus protect against the adverse effects of nutrient overenrichment from cultural eutrophication. EPA's recommended process for developing such criteria includes physical classification of waterbodies, determination of current reference conditions, evaluation of historical data and other information (such as published literature), use of models to simulate physical and ecological processes or determine empirical relationships among causal and response variables (if necessary), expert judgment, and evaluation of downstream effects. EPA has used elements of this process to produce the information contained in this document. The causal (total nitrogen, total phosphorus) and biological and physical response (chlorophyll *a*, Secchi) variables represent a set of starting points for States and Tribes to use in establishing their own criteria.

EPA recommends that States and Tribes establish numerical criteria based on section 304(a) guidance, section 304(a) guidance modified to reflect site-specific conditions, or other scientifically defensible methods. For many pollutants, such as toxic chemicals, EPA expects that section 304(a) guidance will provide an appropriate level of protection without further modification. EPA has also published methods for modifying 304(a) criteria, such as the water

effect ratio, on a site-specific basis where conditions warrant modification to achieve the intended level of protection. For nutrients, however, EPA expects that it will usually be necessary for States and authorized Tribes to be more precise in identifying the nutrient levels that protect aquatic life and recreational uses. This can be achieved through criteria modified to reflect a smaller geographic scale than an Ecoregion, such as a subecoregion, the State or Tribe level, or a specific class of waterbodies. Criteria can be refined by grouping data or performing analyses at these smaller geographic scales. Refinement can also occur through further consideration of other elements such as published literature or models.

EPA expects that the values presented in this document generally represent nutrient levels that protect against the adverse effects of cultural overenrichment and are based on information available to the Agency at the time of this publication. However, States and Tribes should critically evaluate this information in light of the specific uses that need to be protected. For example, more sensitive uses may require more stringent criteria to ensure adequate protection. On the other hand, overly stringent levels of protection against cultural eutrophication may actually fall below the natural load of nutrients for certain waterbodies. In cases such as these, the level of nutrients specified may not be sufficient to support a productive fishery. In the criteria derivation process, it is important to distinguish between the natural load associated with a specific waterbody using historical data and expert judgment and current reference conditions. These elements of the criteria derivation process are best addressed by States and Tribes with access to information and local expertise. Therefore, EPA strongly encourages States and Tribes to use the information contained in this document to develop more refined criteria according to the methods described in EPA's technical guidance manuals for specific waterbody types.

To assist in further refinement of nutrient criteria, EPA has established 10 RTAGs (experts from EPA Regional Offices and States/Tribes). In refining criteria, States and authorized Tribes need to provide documentation of data and analyses, along with a defensible rationale, for any new or revised nutrient criteria they submit to EPA for review and approval. As part of EPA's review of State and Tribal standards, EPA intends to seek assurance from the RTAG that proposed criteria are sufficient to protect uses.

In using the information and recommendations in this document and elsewhere to develop numerical criteria or procedures to translate narrative criteria, EPA encourages States and Tribes to:

- Address both chemical causal variables and early indicator response variables. Causal variables are necessary to protect uses before impairment occurs and to maintain downstream uses. Early response variables are necessary to warn of possible impairment and to integrate the effects of variable and potentially unmeasured nutrient loads.
- Include variables that can be measured to determine if standards are met, and variables that can be related to the ultimate sources of excess nutrients.
- Identify appropriate periods of duration (how long) and frequency (how often) of occurrence in addition to magnitude (how much). EPA does not recommend identifying nutrient concentrations that must be met at all times; rather a seasonal or annual averaging period (e.g., based on weekly or biweekly measurements) is considered appropriate.

However, these central tendency measures should apply each season or each year, except under the most extraordinary conditions (e.g., a 100-year flood).

3.0 AREA COVERED BY THIS DOCUMENT

This chapter provides a general description of the Aggregate Ecoregion and its geographical boundaries. Descriptions of the level III subcoregions contained within the Aggregate Ecoregion are also provided.

3.1 Description of Aggregate Ecoregion V

The nearly level, rolling, and irregular plains of Ecoregion V are mostly semiarid and originally supported mostly grassland. A large part of the area is now cropland and includes the major winter wheat growing area of the United States. The annual freeze-free growing season ranges from 120 days in the northwest to over 240 days in the southeast. Mean annual precipitation is typically erratic and generally decreases from about 28 inches in most eastern areas to about 15 inches in the west. Most of the precipitation occurs during the summer and amounts are usually enough to support dryland farming and rangeland. Intermittent and ephemeral streams are common in drier, western portions and/or where irrigation has lowered the water table.

Ecoregion V is in the transition zone between short and tall grass prairies. Purely tall grass is limited to wetter, easternmost areas whereas short grass is exclusively found in the drier, western part; in between, the potential natural vegetation is a mixture of tall and short grass species. Riparian woodland is usually restricted to wetter eastern areas. Scattered ponderosa pine is found on some ridgetops and bluffs especially in Nebraska. In southwestern Oklahoma and central Texas, shortgrass prairie originally grew with scattered, low shrubs and deciduous trees. The region's characteristic mix of short and tall grass prairie is distinct from the forests of Ecoregion II, the sagebrush, shadscale, and creosote bush of the Xeric West (III), the tall grass prairie of the wetter Corn Belt and Northern Great Plains (VI), and the short grass prairie of the drier Great Plains Grass and Shrublands (IV).

Rocks (including salt formations) and soils have affected the alkalinity, dissolved solid, sulfate, sodium, chloride, and suspended sediment concentrations of streams in Ecoregion V. High sulfate concentrations in stream water occur over broad areas; they are primarily the product of soil leaching and the dissolution of gypsum beds although agricultural compounds such as ammonium sulfate and gypsum also can contribute sulfate to rivers (USGS, 1993).

The landforms, soils, and climate of Ecoregion V are more favorable to agriculture than those of the Great Plains Grass and Shrublands (IV). Today, more than half of Ecoregion V is dryland farmed, about a third is rangeland, and most of the remainder is used for irrigated farming. Ecoregion V is dominated by dryland winter wheat, grain sorghum, and alfalfa farming. Irrigated agriculture occurs in areas with sufficient available ground water and along some reaches of perennial rivers. The number of acres under irrigation have increased rapidly in Ecoregion V and produce silage corn, winter wheat, soybeans, grain sorghum, sugar beets, and other livestock feed and forage crops. The use of commercial fertilizer on cropland increased between 1969 and 1992. Cattle feed lots are found throughout most of the southern and central

parts of the South Central Cultivated Great Plains (V); their density is higher than elsewhere in the United States and the number of cattle fattened at feedlots has increased between 1969 and 1992. Cattle graze much of the agriculturally marginal land and ground water is generally adequate to meet the needs of livestock; rangeland is more common than in Ecoregion VI but much less common than in Ecoregions III and IV which are drier. Overall, the land use mosaic of the South Central Cultivated Great Plains (V) is distinct from that of the surrounding regions; winter wheat and feedlots are far more common than in Ecoregions II, III, IV, VI, or IX.

Agricultural runoff has affected regional water quality. Upward trends in nitrite plus nitrate levels in rivers have mirrored increasing nitrogen fertilizer use (USGS, 1993); in the Platte River of central Nebraska between 1993 and 1995, dissolved nitrate concentrations were highest near cropland and pasture that have had intensive applications of chemical fertilizers. Stream sedimentation from agricultural activity has occurred and has negatively impacted aquatic habitat although no-till cropping practices and other conservation efforts have locally reduced field erosion. Runoff from feedlots is a significant regional water quality problem; nondisinfected runoff from livestock confinement areas have significantly elevated fecal coliform bacteria and nitrite plus nitrate concentrations in rivers (USGS, 1993). Overall, however, potential loss of nitrogen and phosphate fertilizer and pesticide from farm fields is much lower than in the Corn Belt and Northern Great Plains (VI).

Much of the region's irrigation water is obtained from wells or perennial streams. In many areas, ground water withdrawal rates have greatly exceeded recharge rates and have caused the water table to gradually decline, pumping costs to increase, and both springflow and streamflow to diminish. Perennial streams are uncommon and can fluctuate widely in flow from year to year due to the erratic rainfall that characterizes the region. Widespread ground water withdrawal has exacerbated streamflow problems. In addition, irrigation return flows have increased the concentrations of sodium, sulfate, chloride, and dissolved solids in base flows. Irrigation return flows have the greatest impact on river quality during droughts when discharge is dominated by tailwater from irrigated fields. Locally, brines associated with petroleum production have increased the alkalinity, dissolved solid, sulfate, sodium, chloride, and suspended sediment concentrations of streams (USGS, 1993).

Some urban areas in Ecoregion V especially in Texas and Colorado have been growing rapidly and, resultantly, a great deal of agricultural land has been recently converted to urban and industrial uses. Discharges from municipalities and industries have caused elevated concentrations of nitrate and phosphorus in streams. Such increased nutrient loading has caused a corresponding increase in the eutrophication rates of some downstream lakes and reservoirs (USGS, 1993).

3.2 Geographical Boundaries of Aggregate Ecoregion V

Ecoregion V is a fragmented region composed of four separate segments in the central portion of the United States where cultivated Great Plains prevail (Figure 2). The northern most segment begins along the northern border of Montana and goes southeast into the states of North Dakota, South Dakota, and Nebraska.

Aggregate Nutrient Ecoregion 5




Figure 2. Aggregate Ecoregion V.

The second and largest segment is located in the south central portion of the country. It begins in small section of southwestern South Dakota and continues south into Texas. It encompasses the southeastern portion Wyoming, the eastern border of Colorado, and a large portion of Nebraska, Kansas, and Oklahoma. From north central Texas, the segment runs west into eastern New Mexico.

The last two segments are thin strips that run almost parallel from northeast to southeast Texas. The western segment is over double the size of the eastern strip.

3.3 Level III Subcoregions Within Aggregate Ecoregion V

There are four level III subcoregions contained within Aggregate Ecoregion V (Figure 3). The following are brief descriptions provided by Omernik (1999) of the climate, vegetative cover, topography, and other ecological information pertaining to these subcoregions.

25. Western High Plains

Higher and drier than the Central Great Plains to the east, and in contrast to the irregular, mostly grassland or grazing land of the Northwestern Great Plains to the north, much of the Western High Plains comprises smooth to slightly irregular plains having a high percentage of cropland. Grama-buffalo grass is the potential natural vegetation in this region as compared to mostly wheatgrass-needlegrass to the north, Trans-Pecos shrub savanna to the south, and taller grasses to the east. The northern boundary of this ecological region is also the approximate northern limit of winter wheat and sorghum and the southern limit of spring wheat.

27. Central Great Plains

The Central Great Plains are slightly lower, receive more precipitation, and are somewhat more irregular than the Western High Plains to the west. Once a grassland, with scattered low trees and shrubs in the south, much of this ecological region is now cropland, the eastern boundary of the region marking the eastern limits of the major winter wheat growing area of the United States.

32. Texas Blackland Prairies

The Texas Blackland Prairies is a disjunct ecological region distinguished from surrounding regions by its fine textured clayey soils and predominantly prairie potential natural vegetation. This region now contains a higher percent of cropland than adjacent regions, although much of the land has been recently converted to urban and industrial uses.

42. Northwestern Glaciated Plains

The Northwestern Glaciated Plains Ecoregion is a transitional region between the generally more level, moister, more agricultural Northern Glaciated Plains to the east and the generally more irregular, dryer, Northwestern Great Plains to the west and southwest. The western and southwestern boundary roughly coincides with the limits of continental glaciation. Pocking this

Aggregate Nutrient Ecoregion 5 Level III Ecoregions




Figure 3. Aggregate Ecoregion V with level III Ecoregions shown.

Ecoregion is a moderately high concentration of semi-permanent and seasonal wetlands, locally referred to as Prairie Potholes.

3.4 Suggested Ecoregional Subdivisions or Adjustments

EPA recommends that the RTAG evaluate the adequacy of EPA nutrient ecoregional and subecoregional boundaries and refine them as needed to reflect local conditions. See the paper by Dale Robertson (USGS, 2001b) for an alternative approach to Ecoregions entitled “An Alternative Regarding the Scheme for Defining Nutrient Criteria for Rivers and Streams.”

4.0 DATA REVIEW FOR LAKES AND RESERVOIRS IN AGGREGATE ECOREGION V

This section describes the nutrient data EPA has collected and analyzed for this Ecoregion, including an assessment of data quantity and quality. The data tables present the data for each causal parameter (total phosphorus and total nitrogen, both reported and calculated from TKN and nitrite/nitrate) and the primary response variables (Secchi and chlorophyll *a*). EPA considers these parameters essential to nutrient assessment, because the first two are the main causative agents of enrichment and the two response variables are the early indicators of enrichment for most surface waters (see Chapter 5 of the *Lakes and Reservoirs Nutrient Criteria Technical Guidance Manual* [U.S. EPA, 2000a] for a complete discussion on choosing causal and response variables).

4.1 Data Sources

Data sets from Legacy STORET, EPA Region 8 - Montana and Wyoming, EPA Region 8 - South Dakota, and EPA Region 8 - North Dakota were used to assess nutrient conditions from 1990 to 2000. EPA recommends that the RTAGs identify additional data sources that can be used to supplement the data sets listed above. In addition, the RTAGs may utilize published literature values to support quantitative and qualitative analyses.

4.2 Historical Data from Aggregate Ecoregion V (TP, TN, chl *a*, and Secchi)

EPA recommends that States/Tribes assess long-term trends observed over the past 50 years to assess the relative stability of the systems. This information may be obtained from scientific literature or documentation of historical trends. To gain additional perspective on more recent trends, it is recommended that States and Tribes assess nutrient variations over the past 10 years (e.g., what do seasonal variations indicate?).

4.3 QA/QC of Data Sources

An initial quality screen of data was conducted using the rules presented in Appendix C. Data remaining after screening for duplications and other QA measures (e.g., poor or unreported analytical records, sampling errors or omissions, stations associated with outfalls, stormwater sewers, hazardous waste sites) were used in the statistical analyses.

States within Ecoregion V were contacted regarding the quality of their data and information on the methods used to sample and analyze their waters. The following States indicated standard methods or approved EPA methods were used: North Dakota, South Dakota, Montana, Wyoming, Colorado, Kansas, Oklahoma, and Texas. Nebraska indicated that standard or EPA-approved methods were used for some specific nutrient parameters. New Mexico did not provide information prior to the publication of this document.

4.4 Data for All Lakes and Reservoirs Within Aggregate Ecoregion V

Figure 4 shows the location of the sampling stations within each subecoregion. Table 1 presents all data records for all parameters for Aggregate Ecoregion V and subecoregions within the Aggregate Ecoregion.

4.5 Statistical Analysis of Data

EPA's *Technical Guidance Manual for Developing Nutrient Criteria for Lakes and Reservoirs* describes two ways of establishing a reference condition. One method is to choose the upper 25th percentile (75th percentile) of a reference population of lakes. This is the preferred method. The 75th percentile is preferred by EPA because it is likely associated with minimally impacted conditions, will be protective of designated uses, and provides management flexibility. When reference lakes are not identified, the second method is to determine the lower 25th percentile of the population of all lakes within a region to attempt to approximate the preferred approach. The 25th percentile of the entire population was chosen by EPA to represent a surrogate for an actual reference population. Data analyses to date indicate that the lower 25th percentile from an entire population roughly approximates the 75th percentile for a reference population (see case studies for Minnesota lakes in the *Lakes and Reservoirs Nutrient Criteria Technical Guidance Document* [U.S. EPA, 2000a], the case study for Tennessee streams in the *Rivers and Streams Nutrient Criteria Technical Guidance Document* [U.S. EPA, 2000b], the letter from Tennessee Department of Environment and Conservation to Geoffrey Grubbs [TNDEC, 2000], the unpublished paper titled "Estimating the Natural Background Concentrations of Nutrients in Streams and Rivers of the Conterminous United States" [USGS, 2001], and the letter from Matthew Liebman, U.S. EPA Region 1 Nutrient Criteria Coordinator to Geoffrey Grubbs [U.S. EPA, 2000c]). New York State has also presented evidence that the 25th percentile and the 75th percentile compare well based on user perceptions of water resources (NYSDEC, 2000).

Tables 2 and 3a-d present potential reference conditions for both the Aggregate Ecoregion and the subecoregions using both methods. However, the reference lake column is left blank because EPA does not have reference data and anticipates that States/Tribes will provide information on reference lakes. Tables 3a-d present potential reference conditions for rivers and lakes in the level III subecoregions within the Aggregate Ecoregion. Note that the footnotes for Table 2 apply to Tables 3a-d. Appendixes A and B provide a complete presentation of all descriptive statistics for both the Aggregate Ecoregion and the level III subecoregions.

Aggregate Nutrient Ecoregion 5 Lake and Reservoir Stations




Figure 4. Sampling locations within each level III Ecoregion.

Table 1. Lake records* for Aggregate Ecoregion V—South Central Cultivated Great Plains

	Aggregate Ecoregion V	Sub ecoR 25	Sub ecoR 27	Sub ecoR 32	Sub ecoR 42
# of lakes	253	48	116	33	56
# of lake stations	437	75	164	75	123
Key nutrient parameters (listed below)					
- # of records for Secchi depth	1,811	249	463	641	458
- # of records for chlorophyll <i>a</i> (all methods)	805	82	376	336	11
- # of records for total Kjeldhal nitrogen (TKN)	2,644	309	946	790	599
- # of records for nitrite + nitrate (NO ₂ +NO ₃)	1,976	173	708	536	559
- # of records for total nitrogen (TN)	3	3	—	—	—
- # of records for total phosphorus (TP)	3,068	441	1,093	895	639
Total # of records for key nutrient parameters	10,307	1,257	3,586	3,198	2,266

Definitions: (1) # of records refers to the total count of observations for that parameter over the entire decade (1990-1999) for that particular aggregate or subecoregion. These are counts for all seasons over that decade. (2) # of lake stations refers to the total number of lake and reservoir stations within the aggregate or subecoregion from which nutrient data were collected. Since lakes and reservoirs can cross ecoregional boundaries, it is important to note that only those portions of a lake or reservoir (and data associated with those stations) that exist within the Ecoregion are included within this table.

*The number of lakes presented in this table is based on the number of lakes and reservoirs for which nutrient data were provided in the National Nutrient database. This does not imply that this is the total of lakes within the Ecoregion. States and Tribes should determine the representativeness of the tabular data by comparing this information with any additional material they may have.

Table 2. Reference conditions for Aggregate Ecoregion V lakes and reservoirs

Parameter	No. of lakes N*	Reported values		25th percentiles based on all seasons data for the decade	Reference lakes§
		Min	Max	P25† all seasons‡	P75 all seasons
TKN (mg/L)	188	0.13	5.75	0.55	
NO ₂ +NO ₃ -N (mg/L)	162	—	2.70	0.01	
TN (mg/L) - calculated				0.56	
TN (mg/L) - reported	2 (z)	2.42	2.91	2.42 (zz)	
TP (µg/L)	213	1	2,108	33	
Secchi (meters)	171	0.0	4.9	1.3	
Chlorophyll <i>a</i> (µg/L) - F	—	—	—	—	
Chlorophyll <i>a</i> (µg/L) - S	118	0.2	62.2	2.3	
Chlorophyll <i>a</i> (µg/L) - T (u)	2	29	50	29 (zz)	

* N = largest value reported for a decadal season. TN calculated is based on the sum of TKN+NO₂+NO₃. TN reported is actual TN value reported in the database for one sample.

† 75th percentile for Secchi.

‡ Median for all seasons' 25th percentiles, e.g., this value was calculated from four seasons' 25th percentiles. If the seasonal 25th percentile (P25) TP values are: spring 10 µg/L, summer 15 µg/L, fall 12 µg/L, and winter 5 µg/L, the median value of all seasons' P25 will be 11 µg/L.

§ As determined by the Regional Technical Assistance Groups (RTAGs).

Abbreviations: P25, 25th percentile of all data; P75, 75th percentile of all data; F, Chlorophyll *a* measured by Fluorometric method with acid correction; S, Chlorophyll *a* measured by Spectrophotometric method with acid correction; T, Chlorophyll *a* *b c* measured by Trichromatic method; —, not applicable; (u), uncorrected data.

Definitions: (1) Number of Lakes refers to the largest number of lakes and reservoirs for which data existed for a given season within an aggregate nutrient Ecoregion. (2) Medians. All values (min, max, and 25th percentiles) included in the table are based on waterbody medians. All data for a particular parameter within a lake for the decade were reduced to one median for that lake. This prevents over-representation of individual waterbodies with a great deal of data versus those with fewer data points within the statistical analysis. (3) 25th percentile for all seasons is calculated by taking the median of the 4 seasonal 25th percentiles. If a season is missing, the median was calculated with 3 seasons of data. If fewer than 3 seasons were used to derive the median, the entry is flagged (z). (4) A 25th percentile for a season is best derived with data from a minimum of 4 lakes/season. However, this table provides 25th percentiles that were derived with fewer than 4 lakes/season in order to retain all information for all seasons. In calculating the 25th percentile for a season with fewer than 4 lake medians, the statistical program automatically used the minimum value within the fewer-than-4 population. If fewer than 4 lakes were used in developing a seasonal quartile and or all-seasons median, the entry is flagged (zz).

Note: For seasonal values, refer to Appendix A, "Descriptive Statistics Data Tables for Aggregate Ecoregion."

**Table 3a. Reference conditions for Ecoregion V lakes and reservoirs
subcoregion 25**

Parameter	No. of lakes N*	Reported values		25th percentiles based on all seasons data for the decade	Reference lakes§
		Min	Max	P25† all seasons‡	P75 all seasons
TKN (mg/L)	24	0.25	2.22	0.49	1.34
NO ₂ +NO ₃ -N (mg/L)	21	0.00	1.30	0.01	0.34
TN (mg/L) - calculated				0.50	
TN (mg/L) - reported	2 (z)	2.42	2.91	2.42 (zz)	2.91
TP (µg/L)	34	1	1,750	24	150
Secchi (meters)	28	0	3.6	1.5	
Chlorophyll <i>a</i> (µg/L) - F	—	—	—	—	
Chlorophyll <i>a</i> (µg/L) - S	12	0.7	38.4	2.4	25.55
Chlorophyll <i>a</i> (µg/L) - T	2 (z)	29	50	29 (zz)	50.07

**Table 3b. Reference conditions for Ecoregion V lakes and reservoirs
subcoregion 27**

Parameter	No. of lakes N*	Reported values		25th percentiles based on all seasons data for the decade	Reference lakes§
		Min	Max	P25† all seasons‡	P75 all seasons
TKN (mg/L)	92	0.13	2.73	0.61	
NO ₂ +NO ₃ -N (mg/L)	81	—	2.70	0.01	
TN (mg/L) - calculated				0.62	
TN (mg/L) - reported	—	—	—	—	
TP (µg/L)	104	2	570	32	
Secchi (meters)	86	0.1	1.9	0.8	
Chlorophyll <i>a</i> (µg/L) - F	—	—	—	—	
Chlorophyll <i>a</i> (µg/L) - S	78	1.1	56.5	5.8	
Chlorophyll <i>a</i> (µg/L) - T	—	—	—	—	

**Table 3c. Reference conditions for Ecoregion V lakes and reservoirs
subecoregion 32**

Parameter	No. of lakes N*	Reported values		25th percentiles based on all seasons data for the decade	Reference lakes§
		Min	Max	P25† all seasons‡	P75 all seasons
TKN (mg/L)	27	0.27	1.19	0.49	
NO ₂ +NO ₃ -N (mg/L)	15	0.02	0.70	0.12	
TN (mg/L) - calculated				0.61	
TN (mg/L) - reported	—	—	—	—	
TP (µg/L)	28	13	442	25	
Secchi (meters)	29	0.2	2.4	1.1	
Chlorophyll <i>a</i> (µg/L) - F	—	—	—	—	
Chlorophyll <i>a</i> (µg/L) - S	25	0.2	22	1.7	
Chlorophyll <i>a</i> (µg/L) - T	—	—	—	—	

Table 3d. Reference conditions for Ecoregion V lakes and reservoirs subecoregion 42

Parameter	No. of lakes N*	Reported values		25th percentiles based on all seasons data for the decade	Reference lakes§
		Min	Max	P25† all seasons‡	P75 all seasons
TKN (mg/L)	45	0.27	5.75	1.31	
NO ₂ +NO ₃ -N (mg/L)	45	0.00	0.48	0.05	
TN (mg/L) - calculated				1.36	
TN (mg/L) - reported	—	—	—	—	
TP (µg/L)	47	15	1,237	40	
Secchi (meters)	28	0.2	4.4	2.4	
Chlorophyll <i>a</i> (µg/L) - F	—	—	—	—	
Chlorophyll <i>a</i> (µg/L) - S	3 (z)	0.2	9.2	0.2 (zz)	
Chlorophyll <i>a</i> (µg/L) - T	—	—	—	—	

* N = largest value reported for a decadal season. TN calculated is based on the sum of TKN+NO₂+NO₃. TN reported is actual TN value reported in the database for one sample.

† 75th percentile for Secchi.

‡ Median for all seasons' 25th percentiles, e.g., this value was calculated from four seasons' 25th percentiles. If the seasonal 25th percentile (P25) TP values are: spring 10 µg/L, summer 15 µg/L, fall 12 µg/L, and winter 5 µg/L, the median value of all seasons' P25 will be 11 µg/L.

§ As determined by the Regional Technical Assistance Groups (RTAGs).

Abbreviations: P25, 25th percentile of all data; P75, 75th percentile of all data; F, Chlorophyll *a* measured by Fluorometric method with acid correction; S, Chlorophyll *a* measured by Spectrophotometric method with acid correction; T, Chlorophyll *a b c* measured by Trichromatic method; —, not applicable.

Definitions: (1) Number of Lakes refers to the number of lakes and reservoirs for which data existed for the summer months since summer is generally when the greatest amount of nutrient sampling is conducted. If another season greatly predominates, notification is made (s=spring, f=fall, w=winter). (2) Medians. All values (min, max, and 25th percentiles) included in the table are based on waterbody medians. All data for a particular parameter within a lake for the decade were reduced to one median for that lake. This prevents over-representation of individual waterbodies with a great deal of data versus those with fewer data points within the statistical analysis. (3) 25th percentile for all seasons is calculated by taking the median of the 4 seasonal 25th percentiles. If a season is missing, the median was calculated with 3 seasons of data. If fewer than 3 seasons were used to derive the median, the entry is flagged (z). (4) A 25th percentile for a season is best derived with data from a minimum of 4 lakes/season. However, this table provides 25th percentiles that were derived with fewer than 4 lakes/season in order to retain all information for all seasons. In calculating the 25th percentile for a season with fewer than 4 lake medians, the statistical program automatically used the minimum value within the fewer-than-4 population. If fewer than 4 lakes were used in developing a seasonal quartile and or all-seasons median, the entry is flagged (zz).

Note: For seasonal and yearly values, refer to Appendix B, "Descriptive Statistics Data Tables for Level III Subcoregions Within Aggregate Ecoregion."

Table 4 is presented for comparison purposes. It allows the reader to determine where, in the trophic state, the recommended reference conditions fall within traditionally viewed trophic boundaries.

4.6 Classification of Lake/Reservoir Type

Assessing the data by lake type should further reduce the variability in the data analysis. There were no readily available classification data in the national datasets used to develop these criteria. States and Tribes are strongly encouraged to classify their lakes before developing a final criterion.

Table 4. Changes in temperate lake attributes according to trophic state (adapted from Carlson and Simpson, 1995)

TSI Value	SD (m)	TP (µg/L)	Attributes	Water Supply	Recreation	Fisheries
<30	>8	<6	Oligotrophy: Clear water, oxygen throughout the year in the hypolimnion			Salmonid fisheries dominate
30-40	8-4	6-12	Hypolimnia of shallower lakes may become anoxic			Salmonid fisheries in deep lakes
40-50	4-2	12-24	Mesotrophy: Water moderately clear but increasing probability of hypolimnetic anoxia during summer	Iron and manganese evident during the summer. THM precursors exceed 0.1 mg/L and turbidity >1 NTU		Hypolimnetic anoxia results in loss of salmonids. Walleye may predominate
50-60	2-1	24-48	Eutrophy: Anoxic hypolimnia, macrophyte problems possible	Iron, manganese, taste, and odor problems worsen		Warm-water fisheries only. Bass may be dominant
60-70	0.5-1	48-96	Blue-green algae dominate, algal scums and macrophyte problems		Weeds, algal scums, and low transparency discourage swimming and boating	
70-80	0.25-0.5	96-192	Hypereutrophy (light limited). Dense algae and macrophytes			
>80	<0.25	192-384	Algal scums, few macrophytes			Rough fish dominate, summer fish kills possible


Note: This table is provided to allow the reader to make comparisons between the ecoregional criteria provided in this document and traditional nutrient and biological endpoints.

4.7 Summary of Data Reduction Methods

All descriptive statistics were calculated using the medians for each lake within Ecoregion V for which data existed. For example, if one lake had 300 observations for phosphorus over the decade or 1 year's time, one median resulted. Each median from each lake was then used in calculating the percentiles for phosphorus for the aggregate nutrient Ecoregion/subcoregion (level III Ecoregion) by season and year (Figures 5a, 5b).

Preferred Data Choices and Recommendations When Data Are Missing

1. **Where data are missing** or are very low in total records for a given parameter, use 25th percentiles for parameters within an adjacent, similar subcoregion within the same aggregate nutrient Ecoregion, **or** when a similar subcoregion cannot be determined, use the 25th percentile for the Aggregate Ecoregion or consider the **lowest** 25th percentile from a subcoregion (level III) within the aggregate nutrient Ecoregion. Without data, one may assume that the subcoregion in question is as sensitive as the most sensitive subcoregion within the aggregate.
2. **TN calculated:** When reported total nitrogen (TN) median values are lacking or very low in comparison to TKN and Nitrate/Nitrite-N values, the medians for TKN and nitrite/nitrate-N are added, resulting in a calculated TN value. The number of samples (N) for calculated TN is not filled in because it is represented by two subsamples of data: TKN and nitrite/nitrate-N. Therefore, N/A is placed in this box.
3. **TN reported:** This is the median based on reported values for TN from the database.
4. **Chlorophyll *a*:** Medians based on all methods are reported; however, the acid-corrected medians are preferred to the uncorrected medians. In developing a reference condition from a particular method, it is recommended that the method with the most observations be used. Fluorometric and spectrophotometric observations are preferred over all other methods. However, when no data exist for fluorometric and spectrophotometric methods, trichromatic values may be used. Data from the various techniques are not interchangeable.
5. **Periphyton:** Where periphyton data exist, record them separately. For periphyton-dominated streams, a measure of periphyton chlorophyll is a more appropriate response variable than planktonic chlorophyll *a*. See Table 4, page 101, of the *Rivers and Streams Nutrient Technical Guidance Manual* (U.S. EPA, 2000b) for values of periphyton and planktonic chlorophyll *a* related to eutrophy in streams.
6. **Secchi depth:** The 75th percentile is reported for Secchi depth because this is the only variable for which the value of the parameter **increases** with greater clarity (for lakes and reservoirs only).



*Unique Water Body - is a water body that is unique to a state, a subcoregion, a county, the year, and the season.

Figure 5a. Illustration of data reduction process for lake data.




Figure 5b. Illustration of reference condition calculation.

7. **Turbidity units:** Turbidity units from all methods are reported. FTUs and NTUs are preferred over JCUs. If FTUs and NTUs do not exist, use JCUs. These units are not interchangeable. Turbidity is chosen as a response variable in streams because it can be an indicator of increasing algal biomass due to nutrient enrichment. See pages 32-33 of the *Rivers and Streams Nutrient Technical Guidance Manual* for a discussion of turbidity and correlations with algal growth.
8. **Lack of data:** A dash (—) represents missing, inadequate, or inconclusive data. According to EPA statistical analyses, 5% or fewer of the reported observations are “below detection.” Because of this low incidence, these data were retained and factored into the statistical analysis as reported according to the protocols described in Appendix C, “Quality Control/Quality Assurance Rules.”

5.0 REFERENCE SITES AND CONDITIONS IN AGGREGATE ECOREGION V

Reference conditions represent the natural, least impacted conditions, or what is considered to be the most attainable conditions. This chapter compares the different reference conditions determined from the two methods and establishes which reference condition is most appropriate.

- *A priori* determination of reference sites. The preferred method for establishing reference condition is to choose the upper percentile of an *a priori* population of reference lakes. States and Tribes are encouraged to identify reference conditions based on this method.
- Statistical determination of reference conditions (25th percentile of entire database). See Tables 2 and 3a-d in Section 4.0.
- RTAG discussion and rationale for selection of reference sites and conditions in Ecoregion V. The RTAG should compare the results derived from the two methods described above and present a rationale for the final selection of reference sites.

6.0 MODELS USED TO PREDICT OR VERIFY RESPONSE PARAMETERS

The RTAG is encouraged to identify and apply relevant models to support nutrient criteria development. There are three scenarios under which models may be used to derive criteria or support criteria development:

- Models for predicting correlations between causal and response variables
- Models used to verify reference conditions based on percentiles
- Regression models used to predict reference conditions in impacted areas

Appendix C of the Rivers and Streams Technical Guidance Manual (U.S. EPA, 2000b), and Chapter 9 of the Lakes and Reservoirs Technical Guidance Manual (U.S. EPA, 2000a) should be consulted for further details.

7.0 FRAMEWORK FOR REFINING RECOMMENDED NUTRIENT CRITERIA FOR LAKES AND RESERVOIRS IN AGGREGATE ECOREGION V

Information on each of the following six weight-of-evidence factors is important to refine the criteria presented in this document. All elements should be addressed in developing criteria, as is expressed in EPA's nutrient criteria technical guidance manuals. It is our expectation that EPA Regions, States, and Tribes (as RTAGs) will consider these elements as States/Tribes develop their criteria. This section should be viewed as a worksheet (sections are left blank for this purpose) to assist in the refinement of nutrient criteria. If many of these elements are ultimately unaddressed, EPA may rely on the proposed reference conditions presented in Tables 3a-d and other literature and information readily available to the EPA Headquarters nutrient team to develop nutrient water quality recommendations for this Ecoregion.

7.1 Example Worksheet for Developing Aggregate Ecoregion and Subcoregion Nutrient Criteria

Literature sources: _____

Historical data and trends: _____

Reference condition: _____

Models: _____

RTAG expert review and consensus: _____

Downstream effects: _____

7.2 Setting Seasonal Criteria

The recommendations presented in this document are based in part on medians of all the 25th percentile seasonal data (decadal), and as such reflect all seasons and not one particular season or year. It is recommended that States and Tribes monitor in all seasons to best assess compliance with the resulting criterion. States/Tribes may choose to develop criteria that reflect **each** particular season or **given season** or a **given year** when there is significant variability between seasons/years or designated uses that are specifically tied to one or more seasons of the year (e.g., recreation, fishing). Using the tables in Appendix A and B, one can set reference conditions based on a particular season or year and then develop a criterion based on each individual season. Obviously, this option is season-specific and would require increased monitoring within each season to assess compliance. If a case can be made that one season is more appropriate than another season or more appropriate than the annual median, criteria should be season specific. For example, in most parts of the country, spring and summer are the most common growth periods, so criteria for chlorophyll *a* and Secchi may be set for spring and summer only. However, caution should be used when developing criteria for TN and TP because the peak loading of these nutrients may take place in seasons other than summer, such as winter and spring. For these reasons, EPA developed annual criteria and provided additional seasonal information in appendices.

7.3 When Data/Reference Conditions Are Lacking

When data are unavailable to develop a reference condition for a particular parameter(s) within a subcoregion, EPA recommends one of three options: (1) use data from a similar neighboring subcoregion (e.g., if data are few or nonexistent for the Northern Cascades, consider using the data and reference conditions developed for the Cascades); (2) use the 25th percentiles for the Aggregate Ecoregion; or (3) consider using the lowest of the yearly medians for that parameter calculated for all the subcoregions within the Aggregate Ecoregion.

7.4 Site-Specific Criteria Development

Criteria may be refined in a number of ways. The best way is to follow the critical elements of criteria development as well as to refer to the *Lakes and Reservoirs Nutrient Criteria Technical Guidance Manual* (U.S. EPA, 2000a). The Technical Guidance Manual presents sections on each of the following factors to consider in setting criteria:

- Refinements to Ecoregions (Chapter 3). See paper by Dale Robertson (USGS, 2001b), an alternative approach to ecoregions entitled “An Alternative Regarding the Scheme for Defining Nutrient Criteria for Rivers and Streams.”
- Classification of waterbodies (Chapter 3)
- Setting seasonal criteria to reflect major seasonal climate differences and accounting for significant or cyclical precipitation events (high-flow/low-flow conditions) (Chapter 7)
- Setting criteria for reservoirs only. (The technical guidance manual recommends that data be separated for lakes and reservoirs and treated independently if possible because of differing physical conditions that occur in lakes and reservoirs. In this document all data from both reservoirs and lakes were considered together since STORET does not allow for the differentiation of data except by waterbody name.)

8.0 LITERATURE CITED

NYSDEC (New York State Department of Environment and Conservation). 2000. Memorandum from Scott Kishbaugh to Jay Bloomfield, September 26, 2000, regarding reference lakes for nutrient criteria.

Omernik JM. 1999. Primary Distinguishing Characteristics of Level III Ecoregions of the Continental United States. Draft.

Omernik JM. 2000. Draft Aggregations of Level III Ecoregions for the National Nutrient Strategy. [<http://www.epa.gov/ost/standards/ecomap.htm>]

TNDEC (Tennessee Department of Environment and Conservation). 2000. Letter to Geoff Grubbs, October 5, 2000, containing comments on draft nutrient criteria recommendations.

U.S. EPA. 2000a. Nutrient Criteria Technical Guidance Manual: Lakes and Reservoirs. U.S. Environmental Protection Agency, Washington, DC. EPA-822-B00-001.

U.S. EPA. 2000b. Nutrient Criteria Technical Guidance Manual: Rivers and Streams. U.S. Environmental Protection Agency, Washington, DC. EPA-822-B00-002.

U.S. EPA. 2000c. Memorandum from Matthew Liebman to Geoffrey Grubbs, December 15, 2000, regarding comments on draft ambient water quality recommendations for development of numeric nutrient criteria.

USGS (U.S. Geological Survey). 1993. National Water Summary 1990-1991, Water Supply Paper 2400. U.S. Government Printing Office, Washington, DC. 589 pages.

USGS (U.S. Geological Survey). 2001a. Unpublished paper titled: “Estimating the Natural Background Concentrations of Nutrients in Streams and Rivers of the Conterminous United States.” 34 pages.

USGS. 2001b. An Alternative Regarding the Scheme for Defining Nutrient Criteria for Rivers and Streams. Dale M. Robertson, David A. Saad, and Ann Wieben. Water Resources Investigations Report 01-4073.

9.0 APPENDICES

- A. Descriptive Statistics Data Tables for Aggregate Ecoregion
- B. Descriptive Statistics Data Tables for Level III Subcoregions Within Aggregate Ecoregion
- C. Quality Control/Quality Assurance Rules

APPENDIX A

Descriptive Statistics Data Tables for Aggregate Ecoregion

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 1997
Chloro_A_Phyto_Spec_A_u g_L

1

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	25	14.25	.25000	81.10	18.85	3.77	132	0.25	1.79	8.40	18.70	54.30
SPRING	36	10.72	.25000	43.40	10.52	1.75	98	0.25	2.85	7.39	15.25	35.50
SUMMER	118	19.52	.25000	124.45	21.44	1.97	110	2.00	6.55	11.85	25.35	64.35
WINTER	29	7.20	.25000	33.50	9.49	1.76	132	0.25	0.25	2.60	8.01	31.80

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 1993
Chloro_A_Trich_unco_ug_L

2

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
SPRING	2	21.58	.55000	42.60	29.73	21.03	138	0.55	0.55	21.58	42.60	42.60
SUMMER	1	57.53	57.530	57.53	.	.	.	57.53	57.53	57.53	57.53	57.53

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
DIP_ug_L

3

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	5	181.85	16.250	464.50	172.45	77.12	95	16.25	76.00	162.50	190.00	464.50
SPRING	3	55.33	9.5000	137.50	71.32	41.17	129	9.50	9.50	19.00	137.50	137.50
SUMMER	20	403.63	21.500	890.00	271.12	60.62	67	68.25	133.50	393.25	584.25	876.38
WINTER	3	373.17	11.000	1001.00	545.86	315.15	146	11.00	11.00	107.50	1001.0	1001.0

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
 Lakes and Reservoirs
 Descriptive Statistics by Decade and Season
 from 1998 to 1999
 Dissolved_Oxygen_percent_sat

4

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
SUMMER	10	93.26	52.800	164.65	30.06	9.51	32	52.80	74.95	90.35	100.00	164.65

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

5

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	43	8.98	3.5000	14.20	2.10	0.32	23	6.60	7.90	8.60	10.20	12.60
SPRING	51	9.03	.60000	13.50	2.10	0.29	23	6.00	7.90	9.30	9.90	13.15
SUMMER	130	7.28	.02500	12.20	1.83	0.16	25	4.00	6.35	7.19	8.35	10.25
WINTER	61	8.97	.50000	14.13	3.36	0.43	37	1.70	8.60	9.60	11.00	13.00

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

6

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	69	0.15	.00000	2.37	0.32	0.04	213	0.00	0.01	0.06	0.15	0.61
SPRING	78	0.21	.00000	3.03	0.44	0.05	212	0.00	0.01	0.03	0.25	0.82
SUMMER	162	0.14	.00000	3.31	0.33	0.03	242	0.00	0.01	0.02	0.12	0.66
WINTER	81	0.28	.00000	1.87	0.38	0.04	134	0.01	0.03	0.10	0.40	1.00

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

7

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	78	1.02	.12500	6.70	0.92	0.10	90	0.25	0.56	0.78	1.20	2.38
SPRING	96	1.08	.00000	4.69	0.83	0.08	77	0.30	0.54	0.83	1.28	2.90
SUMMER	188	1.21	.12500	4.80	0.81	0.06	67	0.38	0.65	0.91	1.64	2.99
WINTER	86	1.19	.24000	10.60	1.45	0.16	122	0.33	0.45	0.74	1.20	3.37

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
SECCHI_m

8

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	60	1.04	.02500	4.00	0.81	0.10	78	0.19	0.50	0.77	1.45	2.62
SPRING	66	0.95	.02500	5.49	0.96	0.12	101	0.15	0.39	0.60	1.01	2.95
SUMMER	171	0.88	.05080	4.80	0.77	0.06	88	0.18	0.35	0.67	1.19	2.45
WINTER	59	1.25	.02500	5.00	0.91	0.12	73	0.29	0.65	1.02	1.52	3.30

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 1993
Total_Nitrogen_mg_L

9

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
SPRING	1	3.86	3.8600	3.86	.	.	.	3.86	3.86	3.86	3.86	3.86
SUMMER	2	1.47	.97000	1.96	0.70	0.50	48	0.97	0.97	1.47	1.96	1.96

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
Total_Phosphorus_ug_L

10

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	84	145.58	2.5000	2060.00	261.92	28.58	180	8.50	50.00	70.00	127.50	530.50
SPRING	105	114.64	.00000	860.00	165.57	16.16	144	5.00	26.25	55.00	120.00	530.00
SUMMER	213	187.83	.00000	2157.50	289.51	19.84	154	15.00	40.00	70.00	225.00	766.00
WINTER	86	237.80	2.5000	6110.00	738.32	79.62	310	5.00	20.00	40.00	90.00	1355.0

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Decade and Season
from 1990 to 2000
pH_S_U

11

season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
FALL	5	8.45	8.1525	8.82	0.28	0.12	3	8.15	8.21	8.48	8.60	8.82
SPRING	2	8.46	8.4250	8.50	0.05	0.04	1	8.43	8.43	8.46	8.50	8.50
SUMMER	19	8.46	7.9125	8.95	0.29	0.07	3	7.91	8.27	8.48	8.64	8.95
WINTER	3	7.70	6.8700	8.17	0.72	0.42	9	6.87	6.87	8.06	8.17	8.17

Data were not always available for all years.

APPENDIX B

Descriptive Statistics Data tables for Level III Subcoregions Within Aggregate Ecoregion

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subecoregion, Decade and Season
from 1990 to 1997
Chloro_A_Phyto_Spec_A_u_g_L

1

subecoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	2	10.88	2.4500	19.30	11.9	8.43	110	2.45	2.45	10.9	19.3	19.3
25	SPRING	8	16.87	.57000	43.40	15.9	5.61	94	0.57	2.39	15.1	28.0	43.4
25	SUMMER	12	16.60	.75000	54.39	15.4	4.43	92	0.75	5.29	13.7	23.1	54.4
25	WINTER	5	16.98	.25000	33.50	15.8	7.07	93	0.25	1.93	17.4	31.8	33.5
27	FALL	14	22.46	2.1000	81.10	21.5	5.75	96	2.10	8.40	17.9	26.1	81.1
27	SPRING	16	10.05	.25000	31.90	9.08	2.27	90	0.25	3.40	8.35	15.1	31.9
27	SUMMER	78	23.60	2.0000	124.45	24.3	2.75	103	3.05	8.15	15.1	30.8	75.5
27	WINTER	7	3.92	.25000	10.02	4.15	1.57	106	0.25	0.25	2.27	8.01	10.0
32	FALL	7	2.58	.25000	10.90	3.82	1.44	148	0.25	0.25	1.14	3.24	10.9
32	SPRING	12	7.51	.25000	20.00	6.36	1.84	85	0.25	2.54	4.87	13.4	20.0
32	SUMMER	25	9.78	.25000	33.20	7.52	1.50	77	2.00	4.82	8.10	13.2	25.4
32	WINTER	17	5.68	.25000	24.15	7.20	1.75	127	0.25	0.90	2.60	7.60	24.2
42	FALL	2	0.94	.25000	1.63	0.97	0.69	104	0.25	0.25	0.94	1.63	1.63
42	SUMMER	3	6.24	.25000	16.84	9.21	5.32	148	0.25	0.25	1.63	16.8	16.8

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 1993
Chloro_A_Trich_unco_ug_L

2

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	SPRING	2	21.58	.55000	42.60	29.7	21.0	138	0.55	0.55	21.6	42.6	42.6
25	SUMMER	1	57.53	57.530	57.53	.	.	.	57.5	57.5	57.5	57.5	57.5

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
DIP_ug_L

3

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	FALL	5	181.85	16.250	464.50	172	77.1	95	16.3	76.0	163	190	465
42	SPRING	3	55.33	9.5000	137.50	71.3	41.2	129	9.50	9.50	19.0	138	138
42	SUMMER	20	403.63	21.500	890.00	271	60.6	67	68.3	134	393	584	876
42	WINTER	3	373.17	11.000	1001.00	546	315	146	11.0	11.0	108	1001	1001

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1998 to 1999
Dissolved_Oxygen_percent_sat

4

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	SUMMER	10	93.26	52.800	164.65	30.1	9.51	32	52.8	75.0	90.4	100	165

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

5

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	14	8.94	6.6000	11.80	1.54	0.41	17	6.60	8.00	8.85	10.2	11.8
25	SPRING	6	8.88	6.3000	10.60	1.66	0.68	19	6.30	7.80	9.13	10.3	10.6
25	SUMMER	15	7.30	4.0000	10.40	2.01	0.52	28	4.00	5.30	7.60	8.50	10.4
25	WINTER	6	10.71	8.2500	13.00	1.61	0.66	15	8.25	9.70	11.0	11.3	13.0
27	FALL	13	10.58	8.0000	14.20	2.05	0.57	19	8.00	8.80	10.1	12.4	14.2
27	SPRING	17	9.40	6.6750	11.70	1.32	0.32	14	6.68	8.88	9.60	9.90	11.7
27	SUMMER	51	7.87	.02500	12.20	2.07	0.29	26	4.80	6.85	7.50	9.60	10.9
27	WINTER	15	10.50	8.2000	13.70	1.85	0.48	18	8.20	8.80	9.90	12.4	13.7
32	FALL	8	6.80	3.5000	8.15	1.70	0.60	25	3.50	5.95	7.45	7.98	8.15
32	SPRING	18	8.45	5.7000	13.40	1.77	0.42	21	5.70	7.30	8.51	9.35	13.4
32	SUMMER	28	6.33	3.2000	8.60	1.32	0.25	21	3.50	5.60	6.60	7.13	8.15
32	WINTER	19	10.33	8.7000	12.60	0.97	0.22	9	8.70	9.50	10.2	11.0	12.6
42	FALL	8	8.65	6.8300	11.30	1.29	0.46	15	6.83	7.99	8.55	9.01	11.3
42	SPRING	10	9.53	.60000	13.50	3.58	1.13	38	0.60	8.95	9.95	11.0	13.5
42	SUMMER	36	7.18	3.9750	10.10	1.44	0.24	20	4.45	6.10	7.33	8.20	9.80
42	WINTER	21	6.13	.50000	14.13	4.12	0.90	67	0.80	2.00	6.80	9.25	11.0

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

6

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	17	0.07	.00000	0.36	0.10	0.03	155	0.00	0.01	0.01	0.04	0.36
25	SPRING	12	0.39	.02500	1.68	0.47	0.14	121	0.03	0.04	0.28	0.48	1.68
25	SUMMER	21	0.18	.00000	1.50	0.34	0.07	191	0.00	0.01	0.05	0.20	0.49
25	WINTER	10	0.46	.00000	1.10	0.39	0.12	86	0.00	0.02	0.50	0.72	1.10
27	FALL	38	0.21	.00000	2.37	0.41	0.07	202	0.00	0.01	0.07	0.18	0.75
27	SPRING	48	0.16	.00000	3.03	0.49	0.07	308	0.00	0.01	0.01	0.03	0.67
27	SUMMER	81	0.14	.00000	3.31	0.42	0.05	293	0.00	0.01	0.01	0.06	0.68
27	WINTER	36	0.25	.00000	1.87	0.38	0.06	154	0.00	0.03	0.06	0.43	0.91
32	FALL	6	0.12	.01000	0.26	0.11	0.04	92	0.01	0.03	0.09	0.23	0.26
32	SPRING	8	0.33	.05500	0.64	0.18	0.06	56	0.06	0.21	0.30	0.45	0.64
32	SUMMER	15	0.17	.00250	0.75	0.22	0.06	125	0.00	0.00	0.12	0.32	0.75
32	WINTER	14	0.43	.02500	1.19	0.36	0.10	85	0.03	0.20	0.27	0.61	1.19
42	FALL	8	0.08	.00375	0.10	0.03	0.01	43	0.00	0.07	0.10	0.10	0.10
42	SPRING	10	0.11	.02025	0.25	0.06	0.02	58	0.02	0.10	0.10	0.14	0.25
42	SUMMER	45	0.10	.00400	0.70	0.14	0.02	141	0.00	0.01	0.05	0.11	0.40
42	WINTER	21	0.16	.00500	1.55	0.33	0.07	207	0.01	0.03	0.06	0.16	0.32

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

7

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	24	0.96	.25000	2.05	0.54	0.11	56	0.25	0.56	0.82	1.35	1.90
25	SPRING	14	1.07	.00000	3.61	1.00	0.27	93	0.00	0.30	0.80	1.44	3.61
25	SUMMER	24	1.04	.25000	2.40	0.58	0.12	56	0.50	0.64	0.87	1.30	2.25
25	WINTER	12	0.81	.30000	1.50	0.45	0.13	55	0.30	0.41	0.71	1.32	1.50
27	FALL	37	0.94	.12500	2.55	0.56	0.09	59	0.30	0.60	0.80	1.13	2.06
27	SPRING	58	0.95	.00000	2.90	0.50	0.07	53	0.39	0.61	0.90	1.16	1.98
27	SUMMER	92	1.03	.12500	3.05	0.54	0.06	53	0.42	0.70	0.82	1.29	1.99
27	WINTER	37	0.76	.31000	1.16	0.24	0.04	32	0.39	0.52	0.75	0.93	1.10
32	FALL	9	0.62	.26000	0.90	0.20	0.07	32	0.26	0.49	0.63	0.76	0.90
32	SPRING	13	0.63	.30000	1.14	0.24	0.07	39	0.30	0.49	0.54	0.70	1.14
32	SUMMER	27	0.68	.26000	1.44	0.32	0.06	47	0.31	0.49	0.60	0.70	1.34
32	WINTER	17	0.55	.27500	1.25	0.27	0.07	50	0.28	0.41	0.45	0.55	1.25
42	FALL	8	2.06	.24000	6.70	2.30	0.81	112	0.24	0.27	1.27	3.24	6.70
42	SPRING	11	2.28	.34500	4.69	1.34	0.40	59	0.35	1.36	2.41	2.90	4.69
42	SUMMER	45	2.00	.29500	4.80	1.03	0.15	52	0.36	1.38	1.73	2.38	3.96
42	WINTER	20	2.76	.24000	10.60	2.38	0.53	86	0.29	1.26	2.27	3.32	8.20

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
SECCHI_m

8

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	20	1.21	.02500	2.73	0.67	0.15	56	0.23	0.77	1.14	1.54	2.62
25	SPRING	18	1.02	.02500	5.49	1.30	0.31	127	0.03	0.30	0.73	1.00	5.49
25	SUMMER	28	1.13	.10000	3.89	0.98	0.18	86	0.13	0.48	0.93	1.49	3.79
25	WINTER	14	1.17	.02500	3.30	0.82	0.22	70	0.03	0.50	1.11	1.50	3.30
27	FALL	20	0.54	.02540	1.82	0.42	0.09	78	0.11	0.29	0.50	0.57	1.63
27	SPRING	21	0.61	.07620	1.56	0.37	0.08	60	0.15	0.43	0.53	0.81	1.31
27	SUMMER	86	0.66	.05080	2.00	0.47	0.05	71	0.10	0.30	0.54	0.89	1.50
27	WINTER	17	1.24	.43180	2.74	0.51	0.12	41	0.43	0.95	1.14	1.45	2.74
32	FALL	7	0.95	.45000	2.15	0.57	0.21	59	0.45	0.61	0.88	1.05	2.15
32	SPRING	16	0.74	.20000	2.50	0.58	0.14	78	0.20	0.42	0.58	0.76	2.50
32	SUMMER	29	1.00	.22000	2.65	0.70	0.13	70	0.25	0.50	0.80	1.23	2.53
32	WINTER	21	0.98	.20500	2.38	0.56	0.12	57	0.42	0.51	0.94	1.20	1.75
42	FALL	13	1.58	.19500	4.00	1.12	0.31	70	0.20	0.71	1.45	2.29	4.00
42	SPRING	11	1.80	.17500	3.68	1.11	0.34	62	0.18	0.46	2.00	2.53	3.68
42	SUMMER	28	1.18	.25000	4.80	1.13	0.21	95	0.26	0.44	0.83	1.37	4.18
42	WINTER	7	2.28	.28500	5.00	1.84	0.69	81	0.29	0.46	1.52	4.22	5.00

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 1993
Total_Nitrogen_mg_L

9

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	SPRING	1	3.86	3.8600	3.86	.	.	.	3.86	3.86	3.86	3.86	3.86
25	SUMMER	2	1.47	.97000	1.96	0.70	0.50	48	0.97	0.97	1.47	1.96	1.96

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
Total_Phosphorus_ug_L

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	FALL	27	183.00	7.5000	2060.00	397	76.4	217	8.50	40.0	70.0	160	610
25	SPRING	21	105.36	.00000	400.00	117	25.6	111	0.00	20.0	60.0	140	380
25	SUMMER	34	223.86	.00000	2157.50	482	82.6	215	8.00	27.5	40.0	160	1800
25	WINTER	11	158.86	2.5000	1440.00	426	128	268	2.50	5.00	20.0	85.0	1440
27	FALL	38	117.70	2.5000	610.00	125	20.2	106	12.5	50.0	85.0	130	440
27	SPRING	57	73.56	2.5000	530.00	86.3	11.4	117	5.00	25.0	45.0	80.0	225
27	SUMMER	104	125.39	5.0000	700.00	136	13.3	108	20.0	40.0	70.0	153	425
27	WINTER	35	52.50	2.5000	300.00	61.5	10.4	117	5.00	20.0	30.0	70.0	190
32	FALL	10	51.44	2.5000	90.00	30.7	9.69	60	2.50	16.3	60.0	70.0	90.0
32	SPRING	16	156.25	20.000	860.00	257	64.2	164	20.0	47.5	60.0	95.0	860
32	SUMMER	28	91.52	10.000	495.00	115	21.8	126	10.0	20.0	50.0	92.5	363
32	WINTER	19	85.20	16.250	390.00	110	25.2	129	16.3	30.0	50.0	75.0	390
42	FALL	9	255.67	5.0000	865.00	305	102	119	5.00	35.0	70.0	492	865
42	SPRING	11	284.70	26.250	715.00	269	81.2	95	26.3	40.0	217	620	715
42	SUMMER	47	357.30	4.5000	1610.00	355	51.8	99	7.00	60.0	255	534	951
42	WINTER	21	726.06	25.000	6110.00	1368	299	188	27.3	40.0	95.0	780	2050

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Decade and Season
from 1990 to 2000
pH_S_U

11

subcoregion	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	FALL	5	8.45	8.1525	8.82	0.28	0.12	3	8.15	8.21	8.48	8.60	8.82
42	SPRING	2	8.46	8.4250	8.50	0.05	0.04	1	8.43	8.43	8.46	8.50	8.50
42	SUMMER	19	8.46	7.9125	8.95	0.29	0.07	3	7.91	8.27	8.48	8.64	8.95
42	WINTER	3	7.70	6.8700	8.17	0.72	0.42	9	6.87	6.87	8.06	8.17	8.17

Data were not always available for all years.

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 1997
Chloro_A_Phyto_Spec_A_ug_L

1

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	SPRING	3	27.33	2.0000	44.50	22.40	12.93	82	2.00	2.00	35.50	44.50	44.50
25	1990	SUMMER	5	13.22	.95000	33.10	15.55	6.96	118	0.95	1.25	3.80	27.00	33.10
25	1990	WINTER	1	2.88	2.8750	2.88	.	.	.	2.88	2.88	2.88	2.88	2.88
25	1991	FALL	1	19.30	19.300	19.30	.	.	.	19.30	19.30	19.30	19.30	19.30
25	1991	SPRING	2	27.35	12.400	42.30	21.14	14.95	77	12.40	12.40	27.35	42.30	42.30
25	1991	SUMMER	5	10.00	.25000	29.00	12.11	5.42	121	0.25	2.48	3.10	15.15	29.00
25	1991	WINTER	1	2.70	2.7000	2.70	.	.	.	2.70	2.70	2.70	2.70	2.70
25	1992	SPRING	5	21.00	.57000	78.80	33.33	14.91	159	0.57	0.93	4.20	20.50	78.80
25	1992	SUMMER	3	2.32	.25000	5.20	2.57	1.49	111	0.25	0.25	1.50	5.20	5.20
25	1992	WINTER	1	1.93	1.9250	1.93	.	.	.	1.93	1.93	1.93	1.93	1.93
25	1993	FALL	1	2.45	2.4500	2.45	.	.	.	2.45	2.45	2.45	2.45	2.45
25	1993	SPRING	1	0.25	.25000	0.25	.	.	.	0.25	0.25	0.25	0.25	0.25
25	1993	SUMMER	3	25.31	.25000	54.39	27.29	15.76	108	0.25	0.25	21.30	54.39	54.39
25	1993	WINTER	3	13.55	.25000	31.80	16.35	9.44	121	0.25	0.25	8.61	31.80	31.80
25	1994	SUMMER	4	29.98	.25000	111.00	54.15	27.08	181	0.25	0.25	4.33	59.71	111.00
25	1994	WINTER	1	33.50	33.500	33.50	.	.	.	33.50	33.50	33.50	33.50	33.50
25	1995	SUMMER	5	13.56	1.1400	37.80	14.30	6.39	105	1.14	5.20	11.75	11.90	37.80
25	1995	WINTER	1	1.60	1.6000	1.60	.	.	.	1.60	1.60	1.60	1.60	1.60
25	1996	SPRING	3	19.30	14.900	27.70	7.28	4.20	38	14.90	14.90	15.30	27.70	27.70
25	1996	SUMMER	4	10.58	.25000	26.50	11.19	5.60	106	0.25	3.87	7.79	17.30	26.50
25	1996	WINTER	2	13.23	.25000	26.20	18.35	12.98	139	0.25	0.25	13.23	26.20	26.20
25	1997	SUMMER	1	12.30	12.300	12.30	.	.	.	12.30	12.30	12.30	12.30	12.30
27	1990	SUMMER	21	13.77	2.5000	64.95	14.43	3.15	105	3.00	4.90	10.00	13.50	30.65
27	1990	WINTER	2	6.23	.25000	12.20	8.45	5.98	136	0.25	0.25	6.23	12.20	12.20
27	1991	FALL	13	17.95	2.1000	54.30	13.92	3.86	78	2.10	8.40	17.80	18.80	54.30
27	1991	SPRING	4	5.91	1.1250	13.00	5.47	2.74	93	1.13	1.61	4.75	10.20	13.00
27	1991	SUMMER	43	19.99	.25000	88.00	19.41	2.96	97	2.00	8.70	12.90	26.50	58.90
27	1992	FALL	1	10.10	10.100	10.10	.	.	.	10.10	10.10	10.10	10.10	10.10
27	1992	SPRING	7	11.64	3.7000	21.80	7.41	2.80	64	3.70	3.90	10.65	21.40	21.80
27	1992	SUMMER	27	14.86	2.0000	48.25	12.55	2.42	84	3.00	6.00	9.60	19.60	37.00
27	1993	SPRING	3	4.20	2.8000	5.30	1.28	0.74	30	2.80	2.80	4.50	5.30	5.30
27	1993	SUMMER	24	7.93	1.5000	17.90	4.59	0.94	58	1.70	2.75	8.65	10.75	15.00

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 1997
Chloro_A_Phyto_Spec_A ug_L

2

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
27	1994	FALL	2	49.25	17.400	81.10	45.04	31.85	91	17.40	17.40	49.25	81.10	81.10
27	1994	SPRING	11	8.10	.25000	31.90	9.67	2.92	119	0.25	0.25	7.50	7.75	31.90
27	1994	SUMMER	32	28.58	.25000	126.30	34.63	6.12	121	0.25	2.33	12.20	47.60	108.75
27	1995	SUMMER	17	16.45	.25000	45.40	15.88	3.85	96	0.25	4.40	12.10	27.90	45.40
27	1995	WINTER	7	3.61	.25000	10.02	3.72	1.41	103	0.25	0.25	3.81	6.40	10.02
27	1996	SPRING	2	6.27	.25000	12.28	8.51	6.02	136	0.25	0.25	6.27	12.28	12.28
27	1996	SUMMER	8	42.14	1.6000	98.25	31.90	11.28	76	1.60	16.90	40.68	61.08	98.25
27	1997	SUMMER	9	34.91	3.0500	124.45	48.65	16.22	139	3.05	5.20	9.35	37.95	124.45
32	1990	FALL	2	3.63	.25000	7.00	4.77	3.38	132	0.25	0.25	3.63	7.00	7.00
32	1990	SPRING	6	8.43	.50000	20.00	7.15	2.92	85	0.50	1.78	8.14	12.05	20.00
32	1990	SUMMER	21	6.45	.25000	16.60	3.97	0.87	62	1.70	3.45	6.30	8.00	12.30
32	1990	WINTER	3	5.48	.25000	9.20	4.66	2.69	85	0.25	0.25	7.00	9.20	9.20
32	1991	FALL	1	0.50	.50000	0.50	.	.	.	0.50	0.50	0.50	0.50	0.50
32	1991	SPRING	2	9.83	.25000	19.40	13.54	9.58	138	0.25	0.25	9.83	19.40	19.40
32	1991	SUMMER	21	10.21	.25000	32.55	7.35	1.60	72	3.00	4.90	8.90	13.20	22.50
32	1991	WINTER	6	11.90	3.3000	24.15	8.58	3.50	72	3.30	6.00	8.43	21.10	24.15
32	1992	SPRING	5	9.32	.25000	21.80	8.45	3.78	91	0.25	3.80	7.65	13.10	21.80
32	1992	SUMMER	23	13.81	.25000	40.20	10.25	2.14	74	0.25	6.18	15.20	19.20	34.20
32	1992	WINTER	1	0.63	.62500	0.63	.	.	.	0.63	0.63	0.63	0.63	0.63
32	1993	FALL	2	1.75	.25000	3.24	2.11	1.50	121	0.25	0.25	1.75	3.24	3.24
32	1993	SPRING	3	2.81	1.2000	4.42	1.61	0.93	57	1.20	1.20	2.80	4.42	4.42
32	1993	SUMMER	18	12.95	1.0300	33.30	10.77	2.54	83	1.03	3.91	9.06	17.00	33.30
32	1993	WINTER	2	4.70	1.8000	7.60	4.10	2.90	87	1.80	1.80	4.70	7.60	7.60
32	1994	FALL	3	1.28	.25000	3.34	1.78	1.03	139	0.25	0.25	0.25	3.34	3.34
32	1994	SPRING	3	5.91	.25000	15.20	8.11	4.68	137	0.25	0.25	2.28	15.20	15.20
32	1994	SUMMER	14	4.76	.25000	21.50	6.45	1.72	136	0.25	0.25	1.33	8.01	21.50
32	1994	WINTER	6	2.07	.25000	7.50	3.04	1.24	147	0.25	0.25	0.25	3.93	7.50
32	1995	FALL	2	1.14	.25000	2.03	1.26	0.89	110	0.25	0.25	1.14	2.03	2.03
32	1995	SPRING	3	2.67	.25000	5.61	2.72	1.57	102	0.25	0.25	2.16	5.61	5.61
32	1995	SUMMER	11	1.26	.25000	7.69	2.24	0.68	178	0.25	0.25	0.25	1.92	7.69
32	1995	WINTER	8	1.67	.25000	6.41	2.12	0.75	127	0.25	0.25	0.94	2.15	6.41
32	1996	FALL	1	10.90	10.900	10.90	.	.	.	10.90	10.90	10.90	10.90	10.90

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 1997
Chloro_A_Phyto_Spec_A_uq_L

3

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1996	SPRING	3	4.59	3.8400	5.48	0.83	0.48	18	3.84	3.84	4.46	5.48	5.48
32	1996	SUMMER	10	7.78	.25000	32.40	9.38	2.96	121	0.25	1.06	6.52	9.59	32.40
32	1996	WINTER	10	3.49	.25000	16.30	5.03	1.59	144	0.25	0.25	1.04	4.95	16.30
42	1991	SUMMER	1	16.84	16.840	16.84	.	.	.	16.84	16.84	16.84	16.84	16.84
42	1992	FALL	2	0.94	.25000	1.63	0.97	0.69	104	0.25	0.25	0.94	1.63	1.63
42	1992	SUMMER	2	0.94	.25000	1.63	0.97	0.69	104	0.25	0.25	0.94	1.63	1.63

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 1993
Chloro_A_Trich_unco_ug_L

4

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	SPRING	1	42.60	42.600	42.60	.	.	.	42.60	42.60	42.60	42.60	42.60
25	1992	SPRING	1	0.55	.55000	0.55	.	.	.	0.55	0.55	0.55	0.55	0.55
25	1993	SUMMER	1	57.53	57.530	57.53	.	.	.	57.53	57.53	57.53	57.53	57.53

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
DIP_ug_L

5

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1990	SPRING	1	19.00	19.000	19.00	.	.	.	19.00	19.00	19.00	19.00	19.00
42	1990	SUMMER	1	116.50	116.50	116.50	.	.	.	116.50	116.50	116.50	116.50	116.50
42	1991	FALL	4	223.25	76.000	464.50	168.01	84.00	75	76.00	119.25	176.25	327.25	464.50
42	1991	SPRING	1	100.00	100.00	100.00	.	.	.	100.00	100.00	100.00	100.00	100.00
42	1991	SUMMER	15	480.60	64.000	1275.00	388.17	100.23	81	64.00	128.50	424.00	644.00	1275.0
42	1991	WINTER	1	100.00	100.00	100.00	.	.	.	100.00	100.00	100.00	100.00	100.00
42	1992	FALL	1	7.50	7.5000	7.50	.	.	.	7.50	7.50	7.50	7.50	7.50
42	1992	SPRING	1	175.00	175.00	175.00	.	.	.	175.00	175.00	175.00	175.00	175.00
42	1992	SUMMER	2	425.75	49.500	802.00	532.10	376.25	125	49.50	49.50	425.75	802.00	802.00
42	1992	WINTER	2	60.00	5.0000	115.00	77.78	55.00	130	5.00	5.00	60.00	115.00	115.00
42	1993	SPRING	1	9.50	9.5000	9.50	.	.	.	9.50	9.50	9.50	9.50	9.50
42	1993	SUMMER	8	536.44	117.50	965.00	293.84	103.89	55	117.50	294.00	538.00	772.50	965.00
42	1993	WINTER	1	17.00	17.000	17.00	.	.	.	17.00	17.00	17.00	17.00	17.00
42	1994	FALL	1	25.00	25.000	25.00	.	.	.	25.00	25.00	25.00	25.00	25.00
42	1994	SUMMER	9	489.44	21.500	890.00	326.00	108.67	67	21.50	204.50	428.00	768.00	890.00
42	1994	WINTER	1	1001.00	1001.0	1001.00	.	.	.	1001.0	1001.0	1001.0	1001.0	1001.0
42	1995	SUMMER	1	179.50	179.50	179.50	.	.	.	179.50	179.50	179.50	179.50	179.50
42	1998	SUMMER	6	414.50	119.00	844.50	280.07	114.34	68	119.00	138.50	386.25	612.50	844.50
42	1999	SUMMER	4	516.00	226.00	819.00	275.19	137.60	53	226.00	287.75	509.50	744.25	819.00
42	2000	SUMMER	5	208.70	37.000	362.50	127.22	56.90	61	37.00	155.00	189.00	300.00	362.50

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1998 to 1999
Dissolved_Oxygen_percent_sat

6

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1998	SUMMER	6	99.41	52.800	164.65	37.79	15.43	38	52.80	74.95	97.49	109.10	164.65
42	1999	SUMMER	4	84.03	66.200	90.80	11.90	5.95	14	66.20	77.70	89.55	90.35	90.80

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

7

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	FALL	6	8.34	6.6000	10.20	1.38	0.56	17	6.60	7.00	8.58	9.10	10.20
25	1990	SPRING	2	9.55	8.5000	10.60	1.48	1.05	16	8.50	8.50	9.55	10.60	10.60
25	1990	SUMMER	2	7.90	7.3000	8.50	0.85	0.60	11	7.30	7.30	7.90	8.50	8.50
25	1990	WINTER	2	12.00	11.000	13.00	1.41	1.00	12	11.00	11.00	12.00	13.00	13.00
25	1991	FALL	6	9.43	8.0000	11.80	1.52	0.62	16	8.00	8.10	9.05	10.60	11.80
25	1991	SPRING	3	5.67	.10000	8.90	4.84	2.80	85	0.10	0.10	8.00	8.90	8.90
25	1991	SUMMER	11	7.66	4.0000	10.40	2.02	0.61	26	4.00	5.60	7.90	9.20	10.40
25	1991	WINTER	1	11.60	11.600	11.60	.	.	.	11.60	11.60	11.60	11.60	11.60
25	1992	SPRING	4	10.08	8.7000	11.40	1.11	0.56	11	8.70	9.30	10.10	10.85	11.40
25	1992	SUMMER	3	6.87	5.7000	7.60	1.02	0.59	15	5.70	5.70	7.30	7.60	7.60
25	1992	WINTER	1	10.50	10.500	10.50	.	.	.	10.50	10.50	10.50	10.50	10.50
25	1993	FALL	1	7.50	7.5000	7.50	.	.	.	7.50	7.50	7.50	7.50	7.50
25	1993	SPRING	1	9.60	9.6000	9.60	.	.	.	9.60	9.60	9.60	9.60	9.60
25	1993	SUMMER	2	5.00	2.8000	7.20	3.11	2.20	62	2.80	2.80	5.00	7.20	7.20
25	1993	WINTER	2	10.45	9.7000	11.20	1.06	0.75	10	9.70	9.70	10.45	11.20	11.20
25	1994	FALL	1	11.00	11.000	11.00	.	.	.	11.00	11.00	11.00	11.00	11.00
25	1994	SUMMER	1	6.60	6.6000	6.60	.	.	.	6.60	6.60	6.60	6.60	6.60
25	1994	WINTER	1	7.30	7.3000	7.30	.	.	.	7.30	7.30	7.30	7.30	7.30
25	1995	SUMMER	1	9.65	9.6500	9.65	.	.	.	9.65	9.65	9.65	9.65	9.65
25	1995	WINTER	2	9.90	9.2000	10.60	0.99	0.70	10	9.20	9.20	9.90	10.60	10.60
25	1996	SPRING	2	5.75	3.7000	7.80	2.90	2.05	50	3.70	3.70	5.75	7.80	7.80
25	1996	SUMMER	3	3.30	.90000	5.00	2.14	1.23	65	0.90	0.90	4.00	5.00	5.00
25	1997	WINTER	1	11.30	11.300	11.30	.	.	.	11.30	11.30	11.30	11.30	11.30
27	1990	FALL	5	12.36	8.5000	14.40	2.74	1.23	22	8.50	10.40	14.20	14.30	14.40
27	1990	SPRING	11	9.84	6.4500	11.70	1.37	0.41	14	6.45	9.50	9.90	10.90	11.70
27	1990	SUMMER	27	9.10	5.5000	14.00	2.54	0.49	28	6.00	6.75	8.60	11.50	13.20
27	1990	WINTER	3	9.58	9.2500	10.20	0.53	0.31	6	9.25	9.25	9.30	10.20	10.20
27	1991	FALL	9	10.33	8.5000	13.30	1.64	0.55	16	8.50	9.20	10.10	10.80	13.30
27	1991	SPRING	6	7.69	6.4000	9.15	0.90	0.37	12	6.40	7.40	7.55	8.10	9.15
27	1991	SUMMER	23	7.32	5.8000	10.30	1.23	0.26	17	5.90	6.30	7.15	8.00	9.95
27	1991	WINTER	13	10.70	8.2000	13.70	1.93	0.54	18	8.20	8.80	10.70	12.40	13.70
27	1992	SPRING	4	9.45	6.9000	11.90	2.05	1.02	22	6.90	8.13	9.50	10.78	11.90

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

8

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
27	1992	SUMMER	15	7.01	3.8000	10.10	1.45	0.37	21	3.80	6.50	6.90	7.30	10.10
27	1992	WINTER	2	10.03	9.3500	10.70	0.95	0.67	10	9.35	9.35	10.03	10.70	10.70
27	1993	FALL	1	8.00	8.0000	8.00	.	.	.	8.00	8.00	8.00	8.00	8.00
27	1993	SPRING	2	7.80	7.7000	7.90	0.14	0.10	2	7.70	7.70	7.80	7.90	7.90
27	1993	SUMMER	17	7.16	2.9500	8.70	1.28	0.31	18	2.95	7.25	7.40	7.60	8.70
27	1994	FALL	1	9.60	9.6000	9.60	.	.	.	9.60	9.60	9.60	9.60	9.60
27	1994	SUMMER	19	6.99	.02500	10.40	2.22	0.51	32	0.03	6.70	7.20	8.00	10.40
32	1990	FALL	1	6.65	6.6500	6.65	.	.	.	6.65	6.65	6.65	6.65	6.65
32	1990	SPRING	9	8.35	5.1000	9.85	1.58	0.53	19	5.10	7.50	9.30	9.40	9.85
32	1990	SUMMER	16	6.07	2.6000	9.40	1.71	0.43	28	2.60	5.45	6.05	7.15	9.40
32	1990	WINTER	6	9.63	8.1000	10.60	0.89	0.36	9	8.10	9.50	9.55	10.45	10.60
32	1991	FALL	1	9.20	9.2000	9.20	.	.	.	9.20	9.20	9.20	9.20	9.20
32	1991	SPRING	9	7.37	3.8500	13.40	3.01	1.00	41	3.85	5.00	7.75	8.65	13.40
32	1991	SUMMER	18	6.29	2.8000	9.50	1.83	0.43	29	2.80	4.90	6.30	7.40	9.50
32	1991	WINTER	7	9.78	7.5500	11.90	1.73	0.66	18	7.55	8.30	9.60	11.50	11.90
32	1992	FALL	2	6.15	4.9000	7.40	1.77	1.25	29	4.90	4.90	6.15	7.40	7.40
32	1992	SPRING	7	7.88	5.6500	9.70	1.64	0.62	21	5.65	6.20	7.50	9.45	9.70
32	1992	SUMMER	21	5.89	.02500	8.50	2.49	0.54	42	0.10	5.55	6.40	7.40	8.50
32	1992	WINTER	5	9.09	7.9000	10.60	1.14	0.51	13	7.90	8.00	9.30	9.65	10.60
32	1993	FALL	4	5.80	3.5000	7.80	1.96	0.98	34	3.50	4.20	5.95	7.40	7.80
32	1993	SPRING	8	7.93	6.0000	10.05	1.55	0.55	20	6.00	6.35	8.25	9.08	10.05
32	1993	SUMMER	19	7.01	4.0000	11.50	1.73	0.40	25	4.00	6.15	6.90	8.10	11.50
32	1993	WINTER	5	9.69	8.3500	10.70	0.99	0.44	10	8.35	9.20	9.60	10.60	10.70
32	1994	FALL	3	7.93	7.7000	8.10	0.21	0.12	3	7.70	7.70	8.00	8.10	8.10
32	1994	SPRING	5	8.43	7.5000	9.35	0.73	0.33	9	7.50	7.90	8.70	8.70	9.35
32	1994	SUMMER	12	6.34	3.2000	7.25	1.11	0.32	17	3.20	6.30	6.70	7.00	7.25
32	1994	WINTER	8	10.65	9.6000	12.60	1.00	0.35	9	9.60	9.75	10.70	11.05	12.60
32	1995	FALL	2	8.25	8.2000	8.30	0.07	0.05	1	8.20	8.20	8.25	8.30	8.30
32	1995	SPRING	6	8.93	6.7500	10.65	1.34	0.55	15	6.75	8.15	9.25	9.50	10.65
32	1995	SUMMER	11	7.50	4.7000	11.15	1.79	0.54	24	4.70	6.65	7.35	8.50	11.15
32	1995	WINTER	10	10.28	8.7000	11.40	0.83	0.26	8	8.70	9.90	10.40	11.00	11.40
32	1996	FALL	1	7.00	7.0000	7.00	.	.	.	7.00	7.00	7.00	7.00	7.00

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

9

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1996	SPRING	7	8.11	7.0000	9.50	0.93	0.35	12	7.00	7.10	8.15	8.90	9.50
32	1996	SUMMER	11	6.96	5.5000	8.60	0.90	0.27	13	5.50	6.25	7.20	7.30	8.60
32	1996	WINTER	8	10.65	9.3000	11.80	0.75	0.26	7	9.30	10.35	10.55	11.15	11.80
32	1997	SPRING	1	11.80	11.800	11.80	.	.	.	11.80	11.80	11.80	11.80	11.80
32	1997	WINTER	3	11.03	9.5000	12.50	1.50	0.87	14	9.50	9.50	11.10	12.50	12.50
42	1990	FALL	1	9.20	9.2000	9.20	.	.	.	9.20	9.20	9.20	9.20	9.20
42	1990	SPRING	4	10.68	9.4000	13.50	1.91	0.96	18	9.40	9.50	9.90	11.85	13.50
42	1990	SUMMER	2	7.45	5.9000	9.00	2.19	1.55	29	5.90	5.90	7.45	9.00	9.00
42	1990	WINTER	2	1.05	.50000	1.60	0.78	0.55	74	0.50	0.50	1.05	1.60	1.60
42	1991	FALL	4	7.63	6.6000	9.02	1.13	0.56	15	6.60	6.72	7.46	8.55	9.02
42	1991	SPRING	3	9.47	8.1000	10.40	1.21	0.70	13	8.10	8.10	9.90	10.40	10.40
42	1991	SUMMER	27	7.21	3.9750	13.45	2.19	0.42	30	4.14	5.24	7.17	8.65	10.86
42	1991	WINTER	2	5.70	4.8000	6.60	1.27	0.90	22	4.80	4.80	5.70	6.60	6.60
42	1992	FALL	3	11.33	8.8000	14.50	2.90	1.68	26	8.80	8.80	10.70	14.50	14.50
42	1992	SPRING	1	8.00	8.0000	8.00	.	.	.	8.00	8.00	8.00	8.00	8.00
42	1992	SUMMER	4	8.22	7.7800	8.90	0.48	0.24	6	7.78	7.89	8.10	8.55	8.90
42	1992	WINTER	15	6.11	.80000	14.00	4.52	1.17	74	0.80	1.90	6.80	9.00	14.00
42	1993	FALL	2	8.90	8.8000	9.00	0.14	0.10	2	8.80	8.80	8.90	9.00	9.00
42	1993	SPRING	4	8.49	.60000	13.15	5.47	2.74	64	0.60	5.05	10.10	11.93	13.15
42	1993	SUMMER	10	6.62	3.1600	11.44	2.42	0.77	37	3.16	4.69	6.29	8.30	11.44
42	1993	WINTER	3	13.05	11.500	14.25	1.41	0.81	11	11.50	11.50	13.40	14.25	14.25
42	1994	FALL	3	8.73	8.1000	9.30	0.60	0.35	7	8.10	8.10	8.80	9.30	9.30
42	1994	SPRING	2	10.98	10.750	11.20	0.32	0.23	3	10.75	10.75	10.98	11.20	11.20
42	1994	SUMMER	11	5.94	2.2500	10.77	2.43	0.73	41	2.25	3.90	6.30	7.60	10.77
42	1994	WINTER	2	8.75	6.7000	10.80	2.90	2.05	33	6.70	6.70	8.75	10.80	10.80
42	1995	FALL	1	8.30	8.3000	8.30	.	.	.	8.30	8.30	8.30	8.30	8.30
42	1995	SPRING	1	10.90	10.900	10.90	.	.	.	10.90	10.90	10.90	10.90	10.90
42	1995	SUMMER	2	6.78	5.8500	7.70	1.31	0.93	19	5.85	5.85	6.78	7.70	7.70
42	1996	FALL	2	8.70	8.4000	9.00	0.42	0.30	5	8.40	8.40	8.70	9.00	9.00
42	1996	SUMMER	2	6.70	6.4000	7.00	0.42	0.30	6	6.40	6.40	6.70	7.00	7.00
42	1996	WINTER	1	9.25	9.2500	9.25	.	.	.	9.25	9.25	9.25	9.25	9.25
42	1997	WINTER	2	9.60	8.2000	11.00	1.98	1.40	21	8.20	8.20	9.60	11.00	11.00

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Dissolved_Oxygen_mg_L

10

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1998	SUMMER	6	8.01	4.5500	11.75	2.43	0.99	30	4.55	6.36	8.30	8.80	11.75
42	1999	SUMMER	5	7.58	5.7550	9.43	1.33	0.60	18	5.76	7.09	7.78	7.84	9.43
42	2000	SUMMER	5	6.94	5.3600	8.10	1.04	0.46	15	5.36	6.74	6.90	7.60	8.10

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

11

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	FALL	7	0.04	.01000	0.21	0.07	0.03	176	0.01	0.01	0.01	0.04	0.21
25	1990	SPRING	1	0.25	.25000	0.25	.	.	.	0.25	0.25	0.25	0.25	0.25
25	1990	SUMMER	3	0.17	.01000	0.49	0.28	0.16	163	0.01	0.01	0.01	0.49	0.49
25	1990	WINTER	7	0.38	.00000	0.85	0.36	0.14	96	0.00	0.01	0.32	0.72	0.85
25	1991	FALL	7	0.08	.00500	0.31	0.13	0.05	156	0.01	0.01	0.01	0.21	0.31
25	1991	SPRING	1	0.31	.31000	0.31	.	.	.	0.31	0.31	0.31	0.31	0.31
25	1991	SUMMER	8	0.08	.00000	0.38	0.13	0.05	166	0.00	0.00	0.02	0.12	0.38
25	1991	WINTER	1	0.70	.70000	0.70	.	.	.	0.70	0.70	0.70	0.70	0.70
25	1992	FALL	3	0.02	.00250	0.06	0.03	0.02	113	0.00	0.00	0.02	0.06	0.06
25	1992	SPRING	3	0.20	.04000	0.30	0.14	0.08	70	0.04	0.04	0.25	0.30	0.30
25	1992	SUMMER	3	0.03	.00500	0.09	0.05	0.03	147	0.01	0.01	0.01	0.09	0.09
25	1993	FALL	2	0.01	.00500	0.01	0.00	0.00	0	0.01	0.01	0.01	0.01	0.01
25	1993	SUMMER	6	0.35	.00250	1.50	0.57	0.23	165	0.00	0.01	0.16	0.27	1.50
25	1994	FALL	2	0.10	.05000	0.15	0.07	0.05	71	0.05	0.05	0.10	0.15	0.15
25	1994	SPRING	2	0.03	.02500	0.03	0.00	0.00	0	0.03	0.03	0.03	0.03	0.03
25	1994	SUMMER	2	0.05	.00000	0.10	0.07	0.05	141	0.00	0.00	0.05	0.10	0.10
25	1994	WINTER	1	2.13	2.1300	2.13	.	.	.	2.13	2.13	2.13	2.13	2.13
25	1995	FALL	2	0.10	.08000	0.11	0.02	0.02	22	0.08	0.08	0.10	0.11	0.11
25	1995	SUMMER	6	0.06	.00250	0.20	0.08	0.03	117	0.00	0.00	0.04	0.10	0.20
25	1995	WINTER	2	0.04	.01000	0.07	0.04	0.03	106	0.01	0.01	0.04	0.07	0.07
25	1996	FALL	2	0.00	.00000	0.00	0.00	0.00	.	0.00	0.00	0.00	0.00	0.00
25	1996	SPRING	3	0.77	.00000	1.68	0.85	0.49	110	0.00	0.00	0.63	1.68	1.68
25	1996	SUMMER	2	0.12	.00000	0.24	0.17	0.12	141	0.00	0.00	0.12	0.24	0.24
25	1996	WINTER	1	0.02	.02000	0.02	.	.	.	0.02	0.02	0.02	0.02	0.02
25	1997	FALL	3	0.12	.00000	0.36	0.21	0.12	173	0.00	0.00	0.00	0.36	0.36
25	1997	SPRING	3	0.41	.09000	0.82	0.37	0.21	90	0.09	0.09	0.33	0.82	0.82
25	1997	SUMMER	3	0.15	.00000	0.45	0.26	0.15	173	0.00	0.00	0.00	0.45	0.45
25	1997	WINTER	2	0.55	.37000	0.73	0.25	0.18	46	0.37	0.37	0.55	0.73	0.73
27	1990	FALL	13	0.27	.00500	0.71	0.25	0.07	90	0.01	0.06	0.24	0.42	0.71
27	1990	SPRING	10	0.34	.00000	1.46	0.55	0.17	160	0.00	0.00	0.01	0.39	1.46
27	1990	SUMMER	15	0.07	.00500	0.33	0.10	0.03	141	0.01	0.01	0.03	0.08	0.33
27	1990	WINTER	2	0.56	.00500	1.12	0.79	0.56	140	0.01	0.01	0.56	1.12	1.12

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

12

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
27	1991	FALL	21	0.12	.00500	0.49	0.15	0.03	133	0.01	0.01	0.05	0.15	0.49
27	1991	SPRING	25	0.04	.01000	0.56	0.11	0.02	293	0.01	0.01	0.01	0.01	0.18
27	1991	SUMMER	33	0.04	.00000	0.31	0.07	0.01	175	0.01	0.01	0.01	0.02	0.22
27	1991	WINTER	25	0.26	.02500	0.91	0.29	0.06	113	0.03	0.03	0.10	0.45	0.80
27	1992	FALL	3	0.34	.00000	0.88	0.47	0.27	140	0.00	0.00	0.14	0.88	0.88
27	1992	SPRING	7	0.06	.00500	0.35	0.13	0.05	200	0.01	0.01	0.01	0.05	0.35
27	1992	SUMMER	12	0.20	.00500	0.93	0.34	0.10	172	0.01	0.01	0.01	0.35	0.93
27	1992	WINTER	2	0.34	.01000	0.67	0.47	0.33	137	0.01	0.01	0.34	0.67	0.67
27	1993	FALL	1	1.02	1.0200	1.02	.	.	.	1.02	1.02	1.02	1.02	1.02
27	1993	SUMMER	4	0.77	.23000	1.99	0.83	0.42	109	0.23	0.24	0.42	1.29	1.99
27	1994	FALL	2	0.00	.00000	0.00	0.00	0.00	141	0.00	0.00	0.00	0.00	0.00
27	1994	SPRING	1	0.00	.00250	0.00	.	.	.	0.00	0.00	0.00	0.00	0.00
27	1994	SUMMER	14	0.12	.00250	1.08	0.30	0.08	258	0.00	0.00	0.00	0.02	1.08
27	1995	FALL	7	0.13	.00000	0.68	0.25	0.09	187	0.00	0.03	0.03	0.14	0.68
27	1995	SPRING	5	0.06	.02500	0.20	0.08	0.04	130	0.03	0.03	0.03	0.03	0.20
27	1995	SUMMER	12	0.11	.00000	1.04	0.30	0.09	275	0.00	0.00	0.01	0.06	1.04
27	1995	WINTER	7	0.04	.00250	0.07	0.03	0.01	67	0.00	0.01	0.04	0.07	0.07
27	1996	FALL	5	0.25	.00000	0.81	0.34	0.15	135	0.00	0.00	0.12	0.33	0.81
27	1996	SPRING	6	0.16	.00000	0.60	0.24	0.10	145	0.00	0.00	0.07	0.24	0.60
27	1996	SUMMER	5	0.16	.00000	0.75	0.33	0.15	201	0.00	0.00	0.00	0.07	0.75
27	1997	FALL	12	0.28	.00000	2.37	0.66	0.19	239	0.00	0.00	0.08	0.19	2.37
27	1997	SPRING	12	0.32	.00000	3.03	0.86	0.25	268	0.00	0.01	0.03	0.17	3.03
27	1997	SUMMER	12	0.36	.00000	3.31	0.95	0.27	261	0.00	0.00	0.04	0.10	3.31
27	1997	WINTER	4	0.69	.00000	2.62	1.29	0.65	189	0.00	0.00	0.06	1.37	2.62
32	1990	FALL	1	0.03	.02500	0.03	.	.	.	0.03	0.03	0.03	0.03	0.03
32	1990	SPRING	5	0.35	.02500	1.00	0.39	0.17	112	0.03	0.10	0.30	0.30	1.00
32	1990	SUMMER	6	0.17	.02500	0.40	0.15	0.06	88	0.03	0.03	0.15	0.26	0.40
32	1990	WINTER	5	0.42	.02500	0.85	0.38	0.17	92	0.03	0.20	0.20	0.80	0.85
32	1991	FALL	1	0.44	.44000	0.44	.	.	.	0.44	0.44	0.44	0.44	0.44
32	1991	SPRING	7	0.38	.01000	0.98	0.30	0.11	79	0.01	0.24	0.34	0.40	0.98
32	1991	SUMMER	7	0.07	.01000	0.32	0.12	0.05	162	0.01	0.01	0.01	0.15	0.32
32	1991	WINTER	3	0.40	.23500	0.51	0.15	0.08	36	0.24	0.24	0.46	0.51	0.51

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

13

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1992	FALL	1	0.01	.01000	0.01	.	.	.	0.01	0.01	0.01	0.01	0.01
32	1992	SPRING	5	0.44	.19500	0.64	0.17	0.07	38	0.20	0.40	0.43	0.53	0.64
32	1992	SUMMER	5	0.23	.01000	0.49	0.20	0.09	88	0.01	0.10	0.16	0.37	0.49
32	1992	WINTER	4	0.68	.36000	1.20	0.36	0.18	54	0.36	0.45	0.58	0.90	1.20
32	1994	FALL	1	0.26	.26000	0.26	.	.	.	0.26	0.26	0.26	0.26	0.26
32	1994	WINTER	5	0.50	.17500	0.92	0.37	0.17	75	0.18	0.21	0.29	0.88	0.92
32	1995	FALL	2	0.03	.03000	0.04	0.00	0.00	11	0.03	0.03	0.03	0.04	0.04
32	1995	SPRING	3	0.36	.24000	0.50	0.13	0.08	36	0.24	0.24	0.34	0.50	0.50
32	1995	SUMMER	11	0.17	.00250	0.75	0.23	0.07	135	0.00	0.00	0.11	0.32	0.75
32	1995	WINTER	8	0.34	.07000	1.19	0.36	0.13	105	0.07	0.17	0.24	0.33	1.19
32	1996	FALL	1	0.15	.15000	0.15	.	.	.	0.15	0.15	0.15	0.15	0.15
32	1996	WINTER	3	0.11	.03000	0.24	0.11	0.07	103	0.03	0.03	0.06	0.24	0.24
42	1990	FALL	1	0.10	.10000	0.10	.	.	.	0.10	0.10	0.10	0.10	0.10
42	1990	SPRING	3	0.08	.00100	0.13	0.07	0.04	88	0.00	0.00	0.10	0.13	0.13
42	1990	SUMMER	2	0.20	.00750	0.40	0.28	0.20	136	0.01	0.01	0.20	0.40	0.40
42	1990	WINTER	2	0.11	.03000	0.20	0.12	0.08	104	0.03	0.03	0.11	0.20	0.20
42	1991	FALL	4	0.10	.10000	0.10	0.00	0.00	0	0.10	0.10	0.10	0.10	0.10
42	1991	SPRING	3	0.06	.03000	0.10	0.04	0.02	67	0.03	0.03	0.04	0.10	0.10
42	1991	SUMMER	27	0.31	.00300	0.75	0.32	0.06	103	0.00	0.01	0.33	0.65	0.75
42	1991	WINTER	2	0.06	.01000	0.10	0.06	0.05	116	0.01	0.01	0.06	0.10	0.10
42	1992	FALL	3	0.06	.00250	0.10	0.05	0.03	85	0.00	0.00	0.08	0.10	0.10
42	1992	SPRING	1	0.10	.10000	0.10	.	.	.	0.10	0.10	0.10	0.10	0.10
42	1992	SUMMER	14	0.03	.00250	0.12	0.04	0.01	119	0.00	0.01	0.01	0.05	0.12
42	1992	WINTER	15	0.07	.00500	0.20	0.07	0.02	92	0.01	0.02	0.04	0.15	0.20
42	1993	FALL	2	0.07	.00375	0.15	0.10	0.07	134	0.00	0.00	0.07	0.15	0.15
42	1993	SPRING	5	0.12	.00150	0.25	0.09	0.04	77	0.00	0.10	0.10	0.14	0.25
42	1993	SUMMER	10	0.17	.01700	0.70	0.19	0.06	115	0.02	0.10	0.10	0.15	0.70
42	1993	WINTER	3	0.17	.09500	0.32	0.13	0.07	75	0.10	0.10	0.10	0.32	0.32
42	1994	FALL	1	0.10	.10000	0.10	.	.	.	0.10	0.10	0.10	0.10	0.10
42	1994	SPRING	2	0.13	.10000	0.16	0.04	0.03	31	0.10	0.10	0.13	0.16	0.16
42	1994	SUMMER	11	0.09	.00900	0.15	0.03	0.01	37	0.01	0.10	0.10	0.10	0.15
42	1994	WINTER	2	0.87	.18500	1.55	0.97	0.68	111	0.19	0.19	0.87	1.55	1.55

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrite_Nitrate_NO2_NO3_mg_L

14

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1995	FALL	1	0.06	.06000	0.06	.	.	.	0.06	0.06	0.06	0.06	0.06
42	1995	SPRING	1	0.18	.18000	0.18	.	.	.	0.18	0.18	0.18	0.18	0.18
42	1995	SUMMER	2	0.06	.04250	0.09	0.03	0.02	47	0.04	0.04	0.06	0.09	0.09
42	1995	WINTER	1	0.20	.20000	0.20	.	.	.	0.20	0.20	0.20	0.20	0.20
42	1996	FALL	2	0.06	.05000	0.07	0.01	0.01	18	0.05	0.05	0.06	0.07	0.07
42	1996	SUMMER	2	0.05	.04000	0.07	0.02	0.01	34	0.04	0.04	0.05	0.07	0.07
42	1996	WINTER	1	0.11	.11000	0.11	.	.	.	0.11	0.11	0.11	0.11	0.11
42	1998	SUMMER	6	0.13	.05000	0.50	0.18	0.07	136	0.05	0.05	0.05	0.10	0.50
42	1999	SUMMER	4	0.11	.05000	0.18	0.07	0.03	62	0.05	0.05	0.10	0.16	0.18
42	2000	SUMMER	5	0.11	.10000	0.15	0.02	0.01	20	0.10	0.10	0.10	0.10	0.15

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	FALL	7	0.86	.25000	1.80	0.65	0.25	75	0.25	0.25	0.78	1.50	1.80
25	1990	SPRING	1	3.61	3.6100	3.61	.	.	.	3.61	3.61	3.61	3.61	3.61
25	1990	SUMMER	1	0.25	.25000	0.25	.	.	.	0.25	0.25	0.25	0.25	0.25
25	1990	WINTER	7	0.92	.62000	1.50	0.33	0.13	36	0.62	0.70	0.77	1.27	1.50
25	1991	FALL	6	0.75	.50000	0.95	0.18	0.07	24	0.50	0.64	0.73	0.94	0.95
25	1991	SPRING	1	0.80	.80000	0.80	.	.	.	0.80	0.80	0.80	0.80	0.80
25	1991	SUMMER	7	0.74	.12500	1.05	0.34	0.13	46	0.13	0.51	0.85	1.04	1.05
25	1991	WINTER	1	0.30	.30000	0.30	.	.	.	0.30	0.30	0.30	0.30	0.30
25	1992	FALL	3	0.33	.06750	0.52	0.24	0.14	71	0.07	0.07	0.42	0.52	0.52
25	1992	SUMMER	4	1.07	.51500	2.40	0.90	0.45	84	0.52	0.53	0.68	1.60	2.40
25	1993	FALL	3	0.28	.12500	0.60	0.27	0.16	97	0.13	0.13	0.13	0.60	0.60
25	1993	SPRING	1	0.30	.29500	0.30	.	.	.	0.30	0.30	0.30	0.30	0.30
25	1993	SUMMER	5	1.08	.36750	1.92	0.75	0.33	69	0.37	0.56	0.70	1.85	1.92
25	1993	WINTER	3	1.15	.72000	1.39	0.37	0.21	32	0.72	0.72	1.33	1.39	1.39
25	1994	FALL	2	0.74	.64000	0.84	0.14	0.10	19	0.64	0.64	0.74	0.84	0.84
25	1994	SUMMER	4	0.54	.37750	0.78	0.19	0.09	34	0.38	0.39	0.51	0.69	0.78
25	1995	FALL	8	1.35	.59000	2.05	0.52	0.18	38	0.59	0.98	1.35	1.75	2.05
25	1995	SPRING	5	1.34	.70000	2.25	0.69	0.31	52	0.70	0.80	1.05	1.90	2.25
25	1995	SUMMER	12	0.96	.23000	2.00	0.48	0.14	50	0.23	0.63	0.92	1.23	2.00
25	1995	WINTER	1	0.33	.33000	0.33	.	.	.	0.33	0.33	0.33	0.33	0.33
25	1996	FALL	2	0.74	.63000	0.84	0.15	0.11	20	0.63	0.63	0.74	0.84	0.84
25	1996	SPRING	4	0.98	.48000	1.44	0.52	0.26	54	0.48	0.53	1.00	1.43	1.44
25	1996	SUMMER	5	1.00	.61000	2.06	0.61	0.27	62	0.61	0.61	0.72	0.98	2.06
25	1996	WINTER	2	0.90	.38000	1.42	0.74	0.52	82	0.38	0.38	0.90	1.42	1.42
25	1997	FALL	3	0.89	.77000	1.09	0.17	0.10	19	0.77	0.77	0.82	1.09	1.09
25	1997	SPRING	3	0.00	.00000	0.00	0.00	0.00	.	0.00	0.00	0.00	0.00	0.00
25	1997	SUMMER	4	1.18	.61000	2.25	0.76	0.38	65	0.61	0.63	0.93	1.73	2.25
25	1997	WINTER	2	0.00	.00000	0.00	0.00	0.00	.	0.00	0.00	0.00	0.00	0.00
27	1990	FALL	12	0.68	.12500	1.93	0.47	0.14	69	0.13	0.56	0.61	0.68	1.93
27	1990	SPRING	10	0.90	.57000	1.14	0.17	0.05	19	0.57	0.78	0.90	1.00	1.14
27	1990	SUMMER	23	0.79	.12500	1.45	0.30	0.06	38	0.44	0.61	0.80	0.89	1.30
27	1990	WINTER	2	0.68	.65000	0.70	0.04	0.03	5	0.65	0.65	0.68	0.70	0.70

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
27	1991	FALL	21	0.78	.30000	1.50	0.27	0.06	35	0.50	0.60	0.70	0.83	1.20
27	1991	SPRING	24	1.08	.40000	2.90	0.59	0.12	55	0.40	0.70	1.00	1.15	2.50
27	1991	SUMMER	31	0.80	.12500	2.36	0.44	0.08	56	0.13	0.60	0.70	0.97	1.70
27	1991	WINTER	26	0.82	.40000	1.40	0.23	0.05	28	0.50	0.62	0.83	0.98	1.10
27	1992	FALL	4	0.48	.12500	0.70	0.25	0.12	52	0.13	0.32	0.54	0.64	0.70
27	1992	SPRING	7	0.75	.50000	1.00	0.17	0.07	23	0.50	0.60	0.70	0.90	1.00
27	1992	SUMMER	22	0.92	.08250	3.37	0.65	0.14	71	0.33	0.60	0.83	0.90	1.53
27	1992	WINTER	4	0.80	.49500	1.00	0.22	0.11	27	0.50	0.65	0.85	0.95	1.00
27	1993	FALL	2	1.16	1.0500	1.26	0.15	0.11	13	1.05	1.05	1.16	1.26	1.26
27	1993	SPRING	2	0.90	.80000	1.00	0.14	0.10	16	0.80	0.80	0.90	1.00	1.00
27	1993	SUMMER	29	0.90	.37250	2.10	0.39	0.07	43	0.45	0.70	0.80	1.05	1.90
27	1994	FALL	2	1.16	.85000	1.46	0.43	0.30	37	0.85	0.85	1.16	1.46	1.46
27	1994	SPRING	12	0.78	.40000	1.98	0.50	0.15	64	0.40	0.47	0.61	0.77	1.98
27	1994	SUMMER	18	1.14	.30000	3.83	0.90	0.21	78	0.30	0.65	0.90	1.12	3.83
27	1995	FALL	7	1.07	.55000	2.21	0.65	0.25	61	0.55	0.64	0.77	1.78	2.21
27	1995	SPRING	5	0.89	.55000	1.80	0.52	0.23	59	0.55	0.60	0.70	0.78	1.80
27	1995	SUMMER	24	1.25	.38000	2.77	0.70	0.14	56	0.38	0.65	1.17	1.80	2.41
27	1995	WINTER	7	0.65	.31000	1.16	0.34	0.13	51	0.31	0.39	0.52	1.05	1.16
27	1996	FALL	5	0.62	.00000	1.65	0.68	0.30	110	0.00	0.00	0.71	0.72	1.65
27	1996	SPRING	7	0.94	.65000	1.36	0.24	0.09	25	0.65	0.75	0.91	1.07	1.36
27	1996	SUMMER	13	1.08	.40500	1.95	0.43	0.12	40	0.41	0.85	0.95	1.37	1.95
27	1997	FALL	12	1.41	.70000	2.55	0.62	0.18	44	0.70	0.89	1.18	1.98	2.55
27	1997	SPRING	12	1.03	.00000	2.35	0.75	0.22	73	0.00	0.29	1.23	1.35	2.35
27	1997	SUMMER	22	1.07	.02500	2.99	0.76	0.16	71	0.03	0.53	0.99	1.53	2.28
27	1997	WINTER	4	0.68	.00000	1.37	0.56	0.28	83	0.00	0.30	0.68	1.06	1.37
32	1990	FALL	1	0.50	.50000	0.50	.	.	.	0.50	0.50	0.50	0.50	0.50
32	1990	SPRING	6	0.67	.50000	1.10	0.23	0.09	34	0.50	0.50	0.60	0.74	1.10
32	1990	SUMMER	10	0.69	.40000	1.20	0.29	0.09	42	0.40	0.50	0.58	0.80	1.20
32	1990	WINTER	6	0.73	.50000	1.20	0.28	0.11	38	0.50	0.50	0.64	0.90	1.20
32	1991	FALL	1	0.20	.20000	0.20	.	.	.	0.20	0.20	0.20	0.20	0.20
32	1991	SPRING	7	0.73	.40000	1.10	0.24	0.09	33	0.40	0.55	0.70	0.95	1.10
32	1991	SUMMER	11	0.78	.30000	1.50	0.36	0.11	47	0.30	0.55	0.70	1.10	1.50

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

17

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1991	WINTER	6	0.67	.36000	1.60	0.47	0.19	70	0.36	0.42	0.46	0.75	1.60
32	1992	FALL	1	0.70	.70000	0.70	.	.	.	0.70	0.70	0.70	0.70	0.70
32	1992	SPRING	5	0.33	.20000	0.45	0.10	0.04	30	0.20	0.30	0.30	0.40	0.45
32	1992	SUMMER	5	0.42	.20000	0.60	0.18	0.08	42	0.20	0.35	0.35	0.60	0.60
32	1992	WINTER	5	0.45	.30000	0.60	0.11	0.05	25	0.30	0.40	0.44	0.50	0.60
32	1993	FALL	3	0.60	.26000	0.90	0.32	0.19	54	0.26	0.26	0.63	0.90	0.90
32	1993	SPRING	4	0.45	.30000	0.54	0.11	0.05	24	0.30	0.38	0.49	0.53	0.54
32	1993	SUMMER	19	0.63	.30500	1.44	0.33	0.07	52	0.31	0.42	0.50	0.80	1.44
32	1993	WINTER	2	0.47	.35000	0.59	0.17	0.12	36	0.35	0.35	0.47	0.59	0.59
32	1994	FALL	4	0.57	.49000	0.80	0.15	0.08	27	0.49	0.49	0.50	0.66	0.80
32	1994	SPRING	3	0.67	.20000	1.14	0.47	0.27	70	0.20	0.20	0.68	1.14	1.14
32	1994	SUMMER	13	0.58	.21000	1.34	0.30	0.08	52	0.21	0.42	0.56	0.63	1.34
32	1994	WINTER	6	0.46	.38000	0.59	0.08	0.03	18	0.38	0.39	0.44	0.55	0.59
32	1995	FALL	2	0.50	.48000	0.52	0.03	0.02	6	0.48	0.48	0.50	0.52	0.52
32	1995	SPRING	4	0.45	.30000	0.52	0.10	0.05	22	0.30	0.40	0.49	0.50	0.52
32	1995	SUMMER	11	0.93	.26000	4.60	1.24	0.37	133	0.26	0.49	0.54	0.59	4.60
32	1995	WINTER	8	0.50	.24000	1.38	0.36	0.13	72	0.24	0.35	0.41	0.45	1.38
32	1996	FALL	1	0.76	.76000	0.76	.	.	.	0.76	0.76	0.76	0.76	0.76
32	1996	SPRING	3	0.41	.30000	0.46	0.09	0.05	22	0.30	0.30	0.46	0.46	0.46
32	1996	SUMMER	10	0.73	.33000	2.64	0.68	0.22	94	0.33	0.43	0.52	0.57	2.64
32	1996	WINTER	10	0.49	.23000	1.11	0.24	0.07	48	0.23	0.43	0.45	0.50	1.11
42	1990	FALL	1	4.33	4.3300	4.33	.	.	.	4.33	4.33	4.33	4.33	4.33
42	1990	SPRING	4	2.54	1.5400	3.57	0.89	0.44	35	1.54	1.84	2.52	3.24	3.57
42	1990	SUMMER	2	2.37	2.1050	2.63	0.37	0.26	16	2.11	2.11	2.37	2.63	2.63
42	1990	WINTER	4	4.78	1.2000	10.60	4.15	2.07	87	1.20	1.88	3.67	7.69	10.60
42	1991	FALL	4	4.27	1.5350	9.08	3.38	1.69	79	1.54	1.96	3.24	6.59	9.08
42	1991	SPRING	3	2.76	1.8950	3.98	1.09	0.63	39	1.90	1.90	2.41	3.98	3.98
42	1991	SUMMER	27	2.42	1.0750	7.65	1.42	0.27	59	1.24	1.43	1.92	3.13	4.29
42	1991	WINTER	2	5.51	5.0200	6.00	0.69	0.49	13	5.02	5.02	5.51	6.00	6.00
42	1992	FALL	3	0.50	.32000	0.85	0.30	0.17	59	0.32	0.32	0.34	0.85	0.85
42	1992	SPRING	1	5.40	5.3950	5.40	.	.	.	5.40	5.40	5.40	5.40	5.40
42	1992	SUMMER	14	1.54	.26000	4.26	1.15	0.31	74	0.26	0.91	1.29	1.93	4.26

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Nitrogen_Tot_Kjeldhal_mg_L

18

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1992	WINTER	13	2.49	.97000	5.59	1.20	0.33	48	0.97	2.01	2.18	3.11	5.59
42	1993	FALL	2	0.10	.04500	0.16	0.08	0.06	79	0.05	0.05	0.10	0.16	0.16
42	1993	SPRING	5	2.08	.52500	4.04	1.32	0.59	63	0.53	1.36	2.00	2.45	4.04
42	1993	SUMMER	10	1.59	.28000	4.08	1.05	0.33	66	0.28	1.24	1.44	1.73	4.08
42	1993	WINTER	3	0.83	.36000	1.61	0.68	0.39	81	0.36	0.36	0.53	1.61	1.61
42	1994	FALL	1	1.15	1.1500	1.15	.	.	.	1.15	1.15	1.15	1.15	1.15
42	1994	SPRING	2	0.39	.34500	0.44	0.06	0.05	16	0.35	0.35	0.39	0.44	0.44
42	1994	SUMMER	11	1.65	.31000	2.71	0.84	0.25	51	0.31	1.08	1.58	2.39	2.71
42	1994	WINTER	2	2.32	.17250	4.47	3.04	2.15	131	0.17	0.17	2.32	4.47	4.47
42	1995	FALL	1	0.24	.24000	0.24	.	.	.	0.24	0.24	0.24	0.24	0.24
42	1995	SPRING	1	0.25	.24500	0.25	.	.	.	0.25	0.25	0.25	0.25	0.25
42	1995	SUMMER	2	1.00	.36000	1.65	0.91	0.64	91	0.36	0.36	1.00	1.65	1.65
42	1995	WINTER	1	0.33	.33000	0.33	.	.	.	0.33	0.33	0.33	0.33	0.33
42	1996	FALL	2	0.27	.24500	0.29	0.03	0.02	12	0.25	0.25	0.27	0.29	0.29
42	1996	SUMMER	2	0.41	.38500	0.43	0.03	0.02	8	0.39	0.39	0.41	0.43	0.43
42	1996	WINTER	1	0.24	.24000	0.24	.	.	.	0.24	0.24	0.24	0.24	0.24
42	1998	SUMMER	6	1.92	1.3600	2.65	0.52	0.21	27	1.36	1.38	1.88	2.38	2.65
42	1999	SUMMER	4	2.13	1.6450	2.67	0.47	0.24	22	1.65	1.74	2.11	2.52	2.67
42	2000	SUMMER	5	2.36	1.2400	4.80	1.41	0.63	60	1.24	1.61	2.02	2.13	4.80

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
SECCHI_m

19

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	FALL	9	1.48	.50000	2.73	0.78	0.26	53	0.50	0.80	1.45	1.80	2.73
25	1990	SPRING	5	0.60	.20000	0.84	0.25	0.11	41	0.20	0.60	0.61	0.76	0.84
25	1990	SUMMER	9	1.28	.30000	3.89	1.20	0.40	94	0.30	0.48	0.64	2.05	3.89
25	1990	WINTER	8	1.32	.88900	1.88	0.37	0.13	28	0.89	0.96	1.36	1.56	1.88
25	1991	FALL	6	0.93	.63500	1.42	0.29	0.12	31	0.64	0.74	0.84	1.09	1.42
25	1991	SPRING	3	0.67	.60000	0.70	0.06	0.03	9	0.60	0.60	0.70	0.70	0.70
25	1991	SUMMER	13	1.10	.50000	2.30	0.55	0.15	50	0.50	0.70	1.02	1.32	2.30
25	1991	WINTER	1	2.25	2.2500	2.25	.	.	.	2.25	2.25	2.25	2.25	2.25
25	1992	FALL	5	0.86	.43180	1.69	0.51	0.23	60	0.43	0.50	0.66	1.00	1.69
25	1992	SPRING	8	0.79	.28000	1.55	0.53	0.19	67	0.28	0.30	0.69	1.26	1.55
25	1992	SUMMER	6	1.49	.50000	2.54	0.66	0.27	45	0.50	1.30	1.45	1.70	2.54
25	1992	WINTER	1	1.35	1.3500	1.35	.	.	.	1.35	1.35	1.35	1.35	1.35
25	1993	FALL	4	1.94	.90000	4.11	1.49	0.75	77	0.90	0.98	1.36	2.90	4.11
25	1993	SPRING	2	2.90	1.9000	3.90	1.41	1.00	49	1.90	1.90	2.90	3.90	3.90
25	1993	SUMMER	8	1.31	.10000	3.87	1.37	0.49	105	0.10	0.18	0.93	2.15	3.87
25	1993	WINTER	4	1.78	.30000	5.30	2.37	1.18	133	0.30	0.40	0.75	3.15	5.30
25	1994	FALL	4	1.15	.02500	2.01	0.92	0.46	79	0.03	0.41	1.29	1.89	2.01
25	1994	SPRING	3	1.07	.10000	2.40	1.19	0.69	112	0.10	0.10	0.70	2.40	2.40
25	1994	SUMMER	7	1.42	.10000	4.52	1.60	0.60	112	0.10	0.20	0.95	2.39	4.52
25	1994	WINTER	2	0.66	.02500	1.30	0.90	0.64	136	0.03	0.03	0.66	1.30	1.30
25	1995	FALL	3	1.78	1.4224	2.21	0.40	0.23	22	1.42	1.42	1.70	2.21	2.21
25	1995	SPRING	1	0.70	.70000	0.70	.	.	.	0.70	0.70	0.70	0.70	0.70
25	1995	SUMMER	8	1.38	.02500	3.71	1.40	0.50	102	0.03	0.25	0.93	2.47	3.71
25	1995	WINTER	2	0.61	.02500	1.20	0.83	0.59	136	0.03	0.03	0.61	1.20	1.20
25	1996	FALL	2	0.82	.63500	1.00	0.26	0.18	32	0.64	0.64	0.82	1.00	1.00
25	1996	SPRING	5	1.10	.02500	2.95	1.12	0.50	101	0.03	0.50	0.84	1.20	2.95
25	1996	SUMMER	8	0.81	.20000	1.70	0.63	0.22	78	0.20	0.20	0.73	1.38	1.70
25	1996	WINTER	3	0.64	.02500	1.60	0.84	0.49	131	0.03	0.03	0.30	1.60	1.60
25	1997	FALL	4	0.97	.81280	1.22	0.17	0.09	18	0.81	0.86	0.93	1.09	1.22
25	1997	SPRING	4	2.40	.76200	5.49	2.19	1.09	91	0.76	0.85	1.67	3.94	5.49
25	1997	SUMMER	5	1.21	.80000	1.60	0.35	0.16	29	0.80	0.93	1.24	1.50	1.60
25	1997	WINTER	2	0.90	.40000	1.40	0.70	0.50	78	0.40	0.40	0.90	1.40	1.40

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
SECCHI_m

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1998	SPRING	1	1.00	1.0000	1.00	.	.	.	1.00	1.00	1.00	1.00	1.00
25	1998	SUMMER	1	1.55	1.5500	1.55	.	.	.	1.55	1.55	1.55	1.55	1.55
27	1990	FALL	8	0.93	.33020	1.98	0.65	0.23	69	0.33	0.44	0.58	1.55	1.98
27	1990	SPRING	9	0.65	.10160	1.14	0.32	0.11	50	0.10	0.51	0.56	0.84	1.14
27	1990	SUMMER	32	0.68	.10160	1.83	0.44	0.08	65	0.20	0.32	0.56	0.87	1.63
27	1990	WINTER	5	1.15	.30000	1.70	0.57	0.25	49	0.30	0.85	1.40	1.47	1.70
27	1991	FALL	8	0.60	.33020	1.32	0.33	0.12	55	0.33	0.38	0.50	0.69	1.32
27	1991	SPRING	4	0.57	.25000	0.96	0.29	0.15	51	0.25	0.37	0.55	0.78	0.96
27	1991	SUMMER	38	0.71	.11430	1.93	0.45	0.07	64	0.15	0.38	0.64	0.90	1.90
27	1991	WINTER	11	1.30	.76200	2.74	0.55	0.17	42	0.76	0.97	1.14	1.45	2.74
27	1992	FALL	3	0.59	.55000	0.66	0.06	0.04	10	0.55	0.55	0.56	0.66	0.66
27	1992	SPRING	4	0.54	.30000	0.89	0.25	0.13	46	0.30	0.37	0.50	0.72	0.89
27	1992	SUMMER	30	0.76	.07620	1.48	0.40	0.07	52	0.10	0.43	0.78	1.02	1.40
27	1992	WINTER	1	1.03	1.0250	1.03	.	.	.	1.03	1.03	1.03	1.03	1.03
27	1993	FALL	1	0.20	.20320	0.20	.	.	.	0.20	0.20	0.20	0.20	0.20
27	1993	SPRING	3	0.67	.25000	1.50	0.72	0.42	108	0.25	0.25	0.25	1.50	1.50
27	1993	SUMMER	27	0.82	.15000	1.90	0.54	0.10	66	0.19	0.25	0.81	1.21	1.86
27	1994	FALL	2	0.35	.20320	0.51	0.21	0.15	60	0.20	0.20	0.35	0.51	0.51
27	1994	SPRING	1	0.69	.68580	0.69	.	.	.	0.69	0.69	0.69	0.69	0.69
27	1994	SUMMER	25	1.00	.10000	5.00	1.04	0.21	104	0.15	0.40	0.65	1.25	2.40
27	1995	FALL	5	0.39	.20000	0.81	0.27	0.12	67	0.20	0.20	0.25	0.50	0.81
27	1995	SPRING	4	0.38	.15240	0.70	0.23	0.12	62	0.15	0.23	0.33	0.53	0.70
27	1995	SUMMER	16	0.58	.07620	1.50	0.38	0.10	67	0.08	0.25	0.55	0.75	1.50
27	1995	WINTER	1	0.95	.95000	0.95	.	.	.	0.95	0.95	0.95	0.95	0.95
27	1996	FALL	5	0.62	.17780	1.65	0.59	0.27	95	0.18	0.30	0.38	0.61	1.65
27	1996	SPRING	4	0.31	.12700	0.46	0.14	0.07	44	0.13	0.23	0.33	0.39	0.46
27	1996	SUMMER	12	0.66	.17780	2.77	0.73	0.21	110	0.18	0.26	0.43	0.63	2.77
27	1997	FALL	7	0.54	.02540	0.91	0.29	0.11	54	0.03	0.33	0.58	0.71	0.91
27	1997	SPRING	7	1.02	.07620	2.06	0.79	0.30	77	0.08	0.15	0.89	1.98	2.06
27	1997	SUMMER	18	0.84	.05080	2.00	0.63	0.15	75	0.05	0.30	0.69	1.50	2.00
27	1997	WINTER	1	0.43	.43180	0.43	.	.	.	0.43	0.43	0.43	0.43	0.43
32	1990	FALL	2	1.20	.60960	1.80	0.84	0.60	70	0.61	0.61	1.20	1.80	1.80

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
SECCHI_m

21

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1990	SPRING	8	0.74	.45500	1.70	0.41	0.15	56	0.46	0.48	0.62	0.81	1.70
32	1990	SUMMER	22	0.99	.20000	3.30	0.80	0.17	81	0.28	0.33	0.80	1.35	2.60
32	1990	WINTER	7	0.53	.28000	0.80	0.20	0.08	39	0.28	0.33	0.50	0.70	0.80
32	1991	FALL	1	2.00	2.0000	2.00	.	.	.	2.00	2.00	2.00	2.00	2.00
32	1991	SPRING	7	0.53	.12000	1.00	0.32	0.12	60	0.12	0.25	0.50	0.90	1.00
32	1991	SUMMER	23	1.12	.19000	2.80	0.63	0.13	56	0.30	0.66	1.08	1.39	2.13
32	1991	WINTER	8	0.65	.13000	1.40	0.41	0.15	63	0.13	0.34	0.58	0.91	1.40
32	1992	FALL	2	2.05	.85000	3.25	1.70	1.20	83	0.85	0.85	2.05	3.25	3.25
32	1992	SPRING	8	0.67	.20000	1.92	0.53	0.19	79	0.20	0.40	0.58	0.65	1.92
32	1992	SUMMER	24	0.82	.18000	2.00	0.51	0.10	61	0.20	0.40	0.76	1.20	1.80
32	1992	WINTER	5	0.59	.12000	1.42	0.49	0.22	83	0.12	0.43	0.45	0.55	1.42
32	1993	FALL	3	1.22	.45000	2.30	0.96	0.56	79	0.45	0.45	0.90	2.30	2.30
32	1993	SPRING	8	0.69	.12000	1.50	0.50	0.18	73	0.12	0.35	0.56	1.05	1.50
32	1993	SUMMER	21	0.98	.20000	2.80	0.78	0.17	80	0.20	0.50	0.76	1.23	2.65
32	1993	WINTER	4	1.16	.10000	1.93	0.80	0.40	69	0.10	0.55	1.30	1.77	1.93
32	1994	FALL	3	0.79	.63500	0.92	0.14	0.08	18	0.64	0.64	0.81	0.92	0.92
32	1994	SPRING	5	1.27	.33000	2.50	0.96	0.43	76	0.33	0.67	0.75	2.10	2.50
32	1994	SUMMER	17	1.25	.20000	2.80	0.93	0.23	74	0.20	0.41	0.98	1.99	2.80
32	1994	WINTER	8	1.23	.68500	2.10	0.61	0.21	49	0.69	0.74	0.98	1.82	2.10
32	1995	FALL	2	1.10	.92000	1.29	0.26	0.18	23	0.92	0.92	1.10	1.29	1.29
32	1995	SPRING	7	0.90	.30000	2.38	0.68	0.26	76	0.30	0.56	0.73	0.92	2.38
32	1995	SUMMER	13	1.39	.22500	3.05	0.91	0.25	65	0.23	0.78	1.19	1.60	3.05
32	1995	WINTER	8	1.19	.46000	2.00	0.63	0.22	53	0.46	0.73	0.91	1.90	2.00
32	1996	SPRING	7	1.16	.24000	3.05	1.10	0.41	95	0.24	0.38	0.74	2.35	3.05
32	1996	SUMMER	15	1.35	.25000	3.50	0.91	0.23	67	0.25	0.70	1.10	2.01	3.50
32	1996	WINTER	13	1.17	.40000	2.75	0.59	0.16	50	0.40	0.85	1.10	1.30	2.75
32	1997	SPRING	1	0.40	.40000	0.40	.	.	.	0.40	0.40	0.40	0.40	0.40
32	1997	WINTER	3	0.69	.35500	1.07	0.36	0.21	52	0.36	0.36	0.64	1.07	1.07
42	1990	FALL	5	2.39	.24000	4.00	1.55	0.69	65	0.24	1.90	2.00	3.80	4.00
42	1990	SPRING	4	2.58	.30000	5.00	1.96	0.98	76	0.30	1.15	2.50	4.00	5.00
42	1990	SUMMER	6	2.63	.40000	6.90	2.99	1.22	113	0.40	0.50	1.00	6.00	6.90
42	1990	WINTER	1	5.00	5.0000	5.00	.	.	.	5.00	5.00	5.00	5.00	5.00

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
SECCHI_m

22

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1991	FALL	8	1.34	.15000	3.90	1.25	0.44	93	0.15	0.44	0.90	2.00	3.90
42	1991	SPRING	3	1.32	.20000	2.00	0.98	0.56	74	0.20	0.20	1.75	2.00	2.00
42	1991	SUMMER	19	1.25	.15000	5.10	1.19	0.27	96	0.15	0.34	1.02	1.47	5.10
42	1991	WINTER	1	0.19	.19000	0.19	.	.	.	0.19	0.19	0.19	0.19	0.19
42	1992	FALL	6	2.40	.31750	5.10	1.75	0.72	73	0.32	1.37	1.92	3.78	5.10
42	1992	SPRING	4	2.66	.15000	5.40	2.24	1.12	84	0.15	0.95	2.55	4.38	5.40
42	1992	SUMMER	8	2.61	.17780	5.44	2.18	0.77	84	0.18	0.82	1.82	5.00	5.44
42	1992	WINTER	2	0.65	.38000	0.91	0.38	0.27	58	0.38	0.38	0.65	0.91	0.91
42	1993	FALL	4	2.62	.60960	4.75	1.98	0.99	76	0.61	0.95	2.56	4.28	4.75
42	1993	SPRING	8	2.21	.05080	4.00	1.18	0.42	53	0.05	1.71	2.08	3.04	4.00
42	1993	SUMMER	14	1.66	.33000	5.00	1.41	0.38	85	0.33	0.45	1.25	2.79	5.00
42	1993	WINTER	3	2.42	.40640	4.72	2.17	1.25	90	0.41	0.41	2.13	4.72	4.72
42	1994	FALL	5	1.69	.76200	3.00	1.10	0.49	65	0.76	0.81	1.10	2.77	3.00
42	1994	SPRING	5	2.24	.90170	4.00	1.27	0.57	57	0.90	1.30	1.98	3.00	4.00
42	1994	SUMMER	16	1.51	.34000	5.00	1.38	0.34	91	0.34	0.63	1.01	1.88	5.00
42	1994	WINTER	1	4.22	4.2164	4.22	.	.	.	4.22	4.22	4.22	4.22	4.22
42	1995	FALL	4	2.53	1.8500	4.00	0.99	0.50	39	1.85	1.93	2.14	3.14	4.00
42	1995	SPRING	3	2.34	.45720	5.00	2.37	1.37	102	0.46	0.46	1.55	5.00	5.00
42	1995	SUMMER	7	2.13	.70000	3.20	1.09	0.41	51	0.70	1.00	2.82	3.00	3.20
42	1996	FALL	3	1.75	.93980	3.00	1.10	0.63	63	0.94	0.94	1.30	3.00	3.00
42	1996	SPRING	3	3.13	2.4000	4.00	0.81	0.47	26	2.40	2.40	3.00	4.00	4.00
42	1996	SUMMER	7	1.56	.20320	4.60	1.48	0.56	95	0.20	0.55	1.25	1.80	4.60
42	1996	WINTER	2	2.22	1.4478	3.00	1.10	0.78	49	1.45	1.45	2.22	3.00	3.00
42	1997	FALL	1	1.45	1.4500	1.45	.	.	.	1.45	1.45	1.45	1.45	1.45
42	1997	SPRING	2	2.33	2.0000	2.65	0.46	0.33	20	2.00	2.00	2.33	2.65	2.65
42	1997	SUMMER	5	2.48	.90000	4.05	1.47	0.66	59	0.90	1.35	2.10	4.00	4.05
42	1997	WINTER	2	1.55	.50800	2.59	1.47	1.04	95	0.51	0.51	1.55	2.59	2.59
42	1998	SUMMER	6	0.71	.42700	1.07	0.25	0.10	35	0.43	0.49	0.68	0.90	1.07
42	1999	SUMMER	4	0.51	.24000	0.78	0.30	0.15	58	0.24	0.26	0.52	0.77	0.78
42	2000	SUMMER	5	0.76	.25000	1.83	0.66	0.29	86	0.25	0.26	0.55	0.91	1.83

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 1993
Total_Nitrogen_mg_L

23

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	SPRING	1	3.86	3.8600	3.86	.	.	.	3.86	3.86	3.86	3.86	3.86
25	1993	SUMMER	2	1.47	.97000	1.96	0.70	0.50	48	0.97	0.97	1.47	1.96	1.96

Aggregate Nutrient Ecoregion: V

24

Lakes and Reservoirs

Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000

Total_Phosphorus_ug_L

subcoregion	year	season	N	MEAN	MIN	MAX	STDEDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
25	1990	FALL	8	118.56	8.5000	610.00	201.20	71.13	170	8.50	27.50	50.00	87.50	610.00
25	1990	SPRING	4	72.50	10.000	210.00	93.23	46.61	129	10.00	15.00	35.00	130.00	210.00
25	1990	SUMMER	8	151.00	5.0000	900.00	307.06	108.56	203	5.00	6.50	30.00	115.00	900.00
25	1990	WINTER	5	16.00	5.0000	40.00	15.97	7.14	100	5.00	5.00	5.00	25.00	40.00
25	1991	FALL	7	161.43	30.000	830.00	295.26	111.60	183	30.00	40.00	50.00	80.00	830.00
25	1991	SPRING	3	36.67	20.000	50.00	15.28	8.82	42	20.00	20.00	40.00	50.00	50.00
25	1991	SUMMER	11	64.55	5.0000	460.00	131.84	39.75	204	5.00	10.00	30.00	40.00	460.00
25	1991	WINTER	2	15.00	10.000	20.00	7.07	5.00	47	10.00	10.00	15.00	20.00	20.00
25	1992	FALL	4	56.25	5.0000	100.00	39.87	19.93	71	5.00	27.50	60.00	85.00	100.00
25	1992	SPRING	7	132.86	20.000	380.00	124.86	47.19	94	20.00	30.00	110.00	200.00	380.00
25	1992	SUMMER	6	29.79	2.5000	100.00	36.39	14.85	122	2.50	6.25	20.00	30.00	100.00
25	1992	WINTER	1	25.00	25.000	25.00	.	.	.	25.00	25.00	25.00	25.00	25.00
25	1993	FALL	3	29.17	7.5000	40.00	18.76	10.83	64	7.50	7.50	40.00	40.00	40.00
25	1993	SPRING	1	25.00	25.000	25.00	.	.	.	25.00	25.00	25.00	25.00	25.00
25	1993	SUMMER	7	323.93	17.500	1800.00	652.48	246.61	201	17.50	20.00	100.00	130.00	1800.0
25	1993	WINTER	3	60.00	20.000	90.00	36.06	20.82	60	20.00	20.00	70.00	90.00	90.00
25	1994	FALL	3	713.33	40.000	2060.00	1166.2	673.33	163	40.00	40.00	40.00	2060.0	2060.0
25	1994	SPRING	1	400.00	400.00	400.00	.	.	.	400.00	400.00	400.00	400.00	400.00
25	1994	SUMMER	6	396.67	2.5000	2230.00	898.51	366.82	227	2.50	12.50	32.50	70.00	2230.0
25	1994	WINTER	1	1400.00	1400.0	1400.00	.	.	.	1400.0	1400.0	1400.0	1400.0	1400.0
25	1995	FALL	8	165.63	40.000	420.00	121.81	43.07	74	40.00	65.00	167.50	200.00	420.00
25	1995	SPRING	5	101.00	20.000	195.00	74.70	33.41	74	20.00	30.00	120.00	140.00	195.00
25	1995	SUMMER	13	258.56	11.250	2085.00	569.75	158.02	220	11.25	30.00	40.00	160.00	2085.0
25	1995	WINTER	2	750.00	20.000	1480.00	1032.4	730.00	138	20.00	20.00	750.00	1480.0	1480.0
25	1996	FALL	2	95.00	40.000	150.00	77.78	55.00	82	40.00	40.00	95.00	150.00	150.00
25	1996	SPRING	4	80.00	.00000	180.00	81.24	40.62	102	0.00	15.00	70.00	145.00	180.00
25	1996	SUMMER	6	141.67	20.000	330.00	141.62	57.82	100	20.00	40.00	75.00	310.00	330.00
25	1996	WINTER	2	60.00	20.000	100.00	56.57	40.00	94	20.00	20.00	60.00	100.00	100.00
25	1997	FALL	3	56.67	50.000	70.00	11.55	6.67	20	50.00	50.00	50.00	70.00	70.00
25	1997	SPRING	3	0.00	.00000	0.00	0.00	0.00	.	0.00	0.00	0.00	0.00	0.00
25	1997	SUMMER	4	92.50	.00000	220.00	110.57	55.28	120	0.00	0.00	75.00	185.00	220.00
25	1997	WINTER	2	15.00	.00000	30.00	21.21	15.00	141	0.00	0.00	15.00	30.00	30.00

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Total_Phosphorus_ug_L

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
27	1990	FALL	12	62.50	5.0000	190.00	60.70	17.52	97	5.00	5.00	50.00	95.00	190.00
27	1990	SPRING	10	49.50	5.0000	260.00	77.01	24.35	156	5.00	5.00	20.00	50.00	260.00
27	1990	SUMMER	34	74.71	5.0000	240.00	57.46	9.85	77	20.00	40.00	60.00	95.00	220.00
27	1990	WINTER	4	85.00	30.000	140.00	49.33	24.66	58	30.00	45.00	85.00	125.00	140.00
27	1991	FALL	21	66.19	20.000	110.00	26.74	5.83	40	20.00	50.00	60.00	90.00	100.00
27	1991	SPRING	28	57.32	20.000	170.00	39.83	7.53	69	20.00	30.00	40.00	65.00	160.00
27	1991	SUMMER	54	77.13	5.0000	465.00	78.83	10.73	102	20.00	35.00	50.00	90.00	240.00
27	1991	WINTER	24	39.17	5.0000	190.00	48.56	9.91	124	5.00	7.50	20.00	50.00	170.00
27	1992	FALL	4	131.25	60.000	230.00	71.46	35.73	54	60.00	85.00	117.50	177.50	230.00
27	1992	SPRING	9	68.33	10.000	190.00	56.90	18.97	83	10.00	40.00	45.00	90.00	190.00
27	1992	SUMMER	32	112.81	10.000	1180.00	211.86	37.45	188	15.00	27.50	50.00	107.50	370.00
27	1992	WINTER	3	123.33	30.000	300.00	153.08	88.38	124	30.00	30.00	40.00	300.00	300.00
27	1993	FALL	2	355.00	50.000	660.00	431.34	305.00	122	50.00	50.00	355.00	660.00	660.00
27	1993	SPRING	4	67.50	30.000	100.00	33.04	16.52	49	30.00	40.00	70.00	95.00	100.00
27	1993	SUMMER	29	102.24	5.0000	580.00	129.91	24.12	127	20.00	30.00	50.00	130.00	380.00
27	1994	FALL	3	145.00	110.00	185.00	37.75	21.79	26	110.00	110.00	140.00	185.00	185.00
27	1994	SPRING	12	57.60	2.5000	265.00	96.65	27.90	168	2.50	2.50	20.00	38.13	265.00
27	1994	SUMMER	34	149.23	2.5000	575.00	164.16	28.15	110	2.50	30.00	65.00	280.00	535.00
27	1995	FALL	7	102.86	30.000	260.00	83.81	31.68	81	30.00	40.00	60.00	160.00	260.00
27	1995	SPRING	4	132.50	30.000	260.00	100.46	50.23	76	30.00	55.00	120.00	210.00	260.00
27	1995	SUMMER	24	107.97	10.000	375.00	98.95	20.20	92	16.25	30.00	70.00	145.00	310.00
27	1995	WINTER	7	38.21	2.5000	60.00	22.39	8.46	59	2.50	20.00	50.00	60.00	60.00
27	1996	FALL	5	84.00	.00000	230.00	94.50	42.26	112	0.00	0.00	90.00	100.00	230.00
27	1996	SPRING	7	95.00	30.000	200.00	58.81	22.23	62	30.00	60.00	75.00	150.00	200.00
27	1996	SUMMER	13	138.08	30.000	315.00	88.54	24.56	64	30.00	75.00	105.00	155.00	315.00
27	1997	FALL	12	220.83	60.000	610.00	183.86	53.08	83	60.00	95.00	172.50	232.50	610.00
27	1997	SPRING	12	123.85	.00000	530.00	147.77	42.66	119	0.00	13.13	92.50	177.50	530.00
27	1997	SUMMER	22	170.91	20.000	610.00	170.49	36.35	100	30.00	50.00	120.00	185.00	555.00
27	1997	WINTER	4	85.00	30.000	150.00	50.00	25.00	59	30.00	50.00	80.00	120.00	150.00
32	1990	FALL	2	31.25	2.5000	60.00	40.66	28.75	130	2.50	2.50	31.25	60.00	60.00
32	1990	SPRING	9	144.44	30.000	860.00	269.53	89.84	187	30.00	40.00	45.00	70.00	860.00
32	1990	SUMMER	23	127.72	5.0000	730.00	160.43	33.45	126	5.00	30.00	80.00	180.00	362.50

Aggregate Nutrient Ecoregion: V

26

Lakes and Reservoirs

Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000

Total_Phosphorus_ug_L

subcoregion	year	season	N	MEAN	MIN	MAX	STDEVE	STDERR	CV	P5	P25	MEDIAN	P75	P95
32	1990	WINTER	6	74.17	40.000	130.00	36.93	15.08	50	40.00	45.00	60.00	110.00	130.00
32	1991	FALL	1	2.50	2.5000	2.50	.	.	.	2.50	2.50	2.50	2.50	2.50
32	1991	SPRING	9	143.89	25.000	755.00	233.33	77.78	162	25.00	30.00	60.00	120.00	755.00
32	1991	SUMMER	24	79.17	10.000	430.00	93.16	19.02	118	20.00	30.00	40.00	85.00	220.00
32	1991	WINTER	8	148.75	20.000	390.00	150.40	53.17	101	20.00	55.00	70.00	265.00	390.00
32	1992	FALL	1	90.00	90.000	90.00	.	.	.	90.00	90.00	90.00	90.00	90.00
32	1992	SPRING	8	54.38	10.000	150.00	45.15	15.96	83	10.00	25.00	40.00	72.50	150.00
32	1992	SUMMER	24	96.98	2.5000	1025.00	206.78	42.21	213	2.50	20.00	50.00	75.00	250.00
32	1992	WINTER	5	65.50	2.5000	115.00	44.38	19.85	68	2.50	50.00	60.00	100.00	115.00
32	1993	FALL	3	60.00	20.000	90.00	36.06	20.82	60	20.00	20.00	70.00	90.00	90.00
32	1993	SPRING	4	325.00	20.000	1150.00	550.42	275.21	169	20.00	40.00	65.00	610.00	1150.0
32	1993	SUMMER	19	112.24	2.5000	740.00	175.21	40.20	156	2.50	30.00	40.00	100.00	740.00
32	1993	WINTER	2	120.00	50.000	190.00	98.99	70.00	82	50.00	50.00	120.00	190.00	190.00
32	1994	FALL	4	33.44	2.5000	60.00	30.88	15.44	92	2.50	6.88	35.63	60.00	60.00
32	1994	SPRING	3	77.50	2.5000	150.00	73.78	42.60	95	2.50	2.50	80.00	150.00	150.00
32	1994	SUMMER	14	79.38	6.2500	430.00	108.36	28.96	137	6.25	20.00	47.50	80.00	430.00
32	1994	WINTER	6	32.50	10.000	60.00	17.82	7.27	55	10.00	20.00	32.50	40.00	60.00
32	1995	FALL	2	25.00	20.000	30.00	7.07	5.00	28	20.00	20.00	25.00	30.00	30.00
32	1995	SPRING	4	37.50	25.000	60.00	15.55	7.77	41	25.00	27.50	32.50	47.50	60.00
32	1995	SUMMER	11	39.09	10.000	170.00	44.21	13.33	113	10.00	20.00	30.00	35.00	170.00
32	1995	WINTER	7	27.50	2.5000	50.00	15.34	5.80	56	2.50	20.00	30.00	40.00	50.00
32	1996	FALL	1	50.00	50.000	50.00	.	.	.	50.00	50.00	50.00	50.00	50.00
32	1996	SPRING	3	17.50	2.5000	30.00	13.92	8.04	80	2.50	2.50	20.00	30.00	30.00
32	1996	SUMMER	10	52.13	10.000	290.00	84.14	26.61	161	10.00	20.00	27.50	40.00	290.00
32	1996	WINTER	10	41.00	10.000	140.00	36.65	11.59	89	10.00	20.00	30.00	40.00	140.00
42	1990	FALL	2	342.50	85.000	600.00	364.16	257.50	106	85.00	85.00	342.50	600.00	600.00
42	1990	SPRING	4	369.25	217.00	620.00	174.08	87.04	47	217.00	268.50	320.00	470.00	620.00
42	1990	SUMMER	3	289.83	50.000	440.00	209.89	121.18	72	50.00	50.00	379.50	440.00	440.00
42	1990	WINTER	4	2367.50	300.00	6110.00	2596.4	1298.2	110	300.00	655.00	1530.0	4080.0	6110.0
42	1991	FALL	5	477.05	11.250	1130.00	421.73	188.60	88	11.25	222.00	491.50	530.50	1130.0
42	1991	SPRING	3	513.33	375.00	715.00	178.63	103.13	35	375.00	375.00	450.00	715.00	715.00
42	1991	SUMMER	28	513.82	20.000	1610.00	427.34	80.76	83	30.00	130.50	502.50	768.00	1320.0

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
Total_Phosphorus_ug_L

27

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1991	WINTER	2	590.00	400.00	780.00	268.70	190.00	46	400.00	400.00	590.00	780.00	780.00
42	1992	FALL	3	33.00	14.000	70.00	32.05	18.50	97	14.00	14.00	15.00	70.00	70.00
42	1992	SPRING	1	870.00	870.00	870.00	.	.	.	870.00	870.00	870.00	870.00	870.00
42	1992	SUMMER	15	196.30	15.000	932.50	238.74	61.64	122	15.00	35.00	90.00	295.00	932.50
42	1992	WINTER	15	435.87	13.000	1870.00	592.19	152.90	136	13.00	40.00	95.00	745.00	1870.0
42	1993	FALL	3	20.00	5.0000	35.00	15.00	8.66	75	5.00	5.00	20.00	35.00	35.00
42	1993	SPRING	5	183.70	12.500	720.00	301.11	134.66	164	12.50	40.00	56.00	90.00	720.00
42	1993	SUMMER	12	430.25	5.0000	1006.50	353.92	102.17	82	5.00	62.50	389.75	713.75	1006.5
42	1993	WINTER	3	42.17	25.000	60.00	17.51	10.11	42	25.00	25.00	41.50	60.00	60.00
42	1994	FALL	2	55.00	50.000	60.00	7.07	5.00	13	50.00	50.00	55.00	60.00	60.00
42	1994	SPRING	2	40.00	40.000	40.00	0.00	0.00	0	40.00	40.00	40.00	40.00	40.00
42	1994	SUMMER	12	438.08	4.5000	1065.00	396.80	114.55	91	4.50	36.75	382.00	829.50	1065.0
42	1994	WINTER	2	592.50	20.000	1165.00	809.64	572.50	137	20.00	20.00	592.50	1165.0	1165.0
42	1995	FALL	1	70.00	70.000	70.00	.	.	.	70.00	70.00	70.00	70.00	70.00
42	1995	SPRING	1	22.25	22.250	22.25	.	.	.	22.25	22.25	22.25	22.25	22.25
42	1995	SUMMER	2	430.25	4.5000	856.00	602.10	425.75	140	4.50	4.50	430.25	856.00	856.00
42	1995	WINTER	1	27.25	27.250	27.25	.	.	.	27.25	27.25	27.25	27.25	27.25
42	1996	FALL	2	27.25	4.5000	50.00	32.17	22.75	118	4.50	4.50	27.25	50.00	50.00
42	1996	SUMMER	3	43.17	4.5000	70.00	34.32	19.81	79	4.50	4.50	55.00	70.00	70.00
42	1996	WINTER	1	30.00	30.000	30.00	.	.	.	30.00	30.00	30.00	30.00	30.00
42	1997	FALL	1	35.00	35.000	35.00	.	.	.	35.00	35.00	35.00	35.00	35.00
42	1997	SUMMER	1	180.00	180.00	180.00	.	.	.	180.00	180.00	180.00	180.00	180.00
42	1998	SUMMER	6	523.42	234.50	970.00	293.91	119.99	56	234.50	249.00	455.50	776.00	970.00
42	1999	SUMMER	4	673.13	369.00	1105.00	346.93	173.47	52	369.00	393.00	609.25	953.25	1105.0
42	2000	SUMMER	5	337.00	224.00	533.50	124.37	55.62	37	224.00	248.00	304.00	375.50	533.50

Aggregate Nutrient Ecoregion: V
Lakes and Reservoirs
Descriptive Statistics by Subcoregion, Year and Season
from 1990 to 2000
pH_S_U

28

subcoregion	year	season	N	MEAN	MIN	MAX	STDDEV	STDERR	CV	P5	P25	MEDIAN	P75	P95
42	1990	FALL	1	8.50	8.5000	8.50	.	.	.	8.50	8.50	8.50	8.50	8.50
42	1990	SPRING	1	8.50	8.5000	8.50	.	.	.	8.50	8.50	8.50	8.50	8.50
42	1990	SUMMER	1	8.90	8.8950	8.90	.	.	.	8.90	8.90	8.90	8.90	8.90
42	1990	WINTER	1	8.17	8.1700	8.17	.	.	.	8.17	8.17	8.17	8.17	8.17
42	1991	FALL	4	8.52	8.2100	8.82	0.26	0.13	3	8.21	8.33	8.53	8.71	8.82
42	1991	SPRING	1	8.50	8.5000	8.50	.	.	.	8.50	8.50	8.50	8.50	8.50
42	1991	SUMMER	15	8.55	7.9500	9.02	0.35	0.09	4	7.95	8.38	8.57	8.80	9.02
42	1991	WINTER	1	7.74	7.7350	7.74	.	.	.	7.74	7.74	7.74	7.74	7.74
42	1992	FALL	1	7.85	7.8500	7.85	.	.	.	7.85	7.85	7.85	7.85	7.85
42	1992	SPRING	1	8.35	8.3500	8.35	.	.	.	8.35	8.35	8.35	8.35	8.35
42	1992	SUMMER	2	8.52	8.3400	8.71	0.26	0.18	3	8.34	8.34	8.52	8.71	8.71
42	1992	WINTER	1	8.39	8.3850	8.39	.	.	.	8.39	8.39	8.39	8.39	8.39
42	1993	SUMMER	8	8.06	7.2750	8.74	0.44	0.16	5	7.28	7.93	7.99	8.31	8.74
42	1994	FALL	1	8.46	8.4550	8.46	.	.	.	8.46	8.46	8.46	8.46	8.46
42	1994	SUMMER	8	8.35	8.0900	8.88	0.28	0.10	3	8.09	8.15	8.27	8.48	8.88
42	1994	WINTER	1	6.87	6.8700	6.87	.	.	.	6.87	6.87	6.87	6.87	6.87
42	1995	SUMMER	1	8.43	8.4300	8.43	.	.	.	8.43	8.43	8.43	8.43	8.43
42	1998	SUMMER	6	8.51	8.2575	8.77	0.21	0.09	3	8.26	8.31	8.51	8.73	8.77
42	1999	SUMMER	4	8.58	8.1770	8.90	0.30	0.15	4	8.18	8.36	8.61	8.79	8.90
42	2000	SUMMER	4	8.32	8.0800	8.55	0.23	0.12	3	8.08	8.12	8.32	8.52	8.55

APPENDIX C

Quality Control/Quality Assurance Rules



Continued Support for the Compilation and Analysis of National Nutrient Data

9 Nutrient Ecoregion/Waterbody Type Summary Chapters

Prepared for:

Steve Potts
Environmental Protection Agency
OW/OST/HECD

Prepared by:

INDUS Corporation
1953 Gallows Road
Vienna, Virginia 22182

Contract Number:68-C-99-226
Task Number:07
Subtask Number:4

August 27, 2001

CONTENTS

1.0	BACKGROUND	C-1
1.1	Purpose	C-1
1.2	References	C-1
2.0	QA/QC PROCEDURES	C-1
2.1	National Data Sets	C-3
2.2	State Data	C-3
2.3	Laboratory Methods	C-4
2.4	Waterbody Name and Class Information	C-4
2.5	Ecoregion Data	C-5
3.0	STATISTICAL ANALYSIS REPORTS	C-5
3.1	Data Source Reports	C-6
3.2	Remark Code Reports	C-6
3.3	Median of Each Waterbody	C-7
3.4	Descriptive Statistic Reports	C-7
3.5	Regression Models	C-7
4.0	TIME PERIOD	C-8
5.0	DATA SOURCES AND PARAMETERS FOR THE AGGREGATE NUTRIENT ECOREGIONS	C-8
5.1	Lakes and Reservoirs	C-9
5.1.1	Aggregate Nutrient Ecoregion 3	C-9
5.1.2	Aggregate Nutrient Ecoregion 4	C-9
5.1.3	Aggregate Nutrient Ecoregion 5	C-10
5.1.4	Aggregate Nutrient Ecoregion 14	C-10
5.2	Rivers and Streams	C-11
5.2.1	Aggregate Nutrient Ecoregion 1	C-11
5.2.2	Aggregate Nutrient Ecoregion 4	C-12
5.2.3	Aggregate Nutrient Ecoregion 5	C-13
5.2.4	Aggregate Nutrient Ecoregion 8	C-13
5.2.5	Aggregate Nutrient Ecoregion 10	C-14
APPENDIX A	Process Used to QA/QA the Legacy STORET Nutrient Data Set	C-16
APPENDIX B	Process for Adding Aggregate Nutrient Ecoregions and Level III Ecoregions	C-22
APPENDIX C	Glossary	C-23

1.0 BACKGROUND

The Nutrient Criteria Program initiated the development of a national Nutrient Criteria Database application that is used to store and analyze nutrient data. The ultimate use of these data is to derive ecoregion specific nutrient criteria. EPA converted STORage and RETrieval (STORET) legacy data, National Stream Quality Accounting Network (NASQAN) data, National Water-Quality Assessment (NAWQA) data, and other relevant nutrient data from universities and States/Tribes into the database. The data imported into the Nutrient Criteria Database are used to develop national nutrient criteria recommendations.

1.1 Purpose

The purpose of this deliverable is to provide EPA with information regarding the database used to create the statistical reports which will be used to derive ecoregion-specific nutrient criteria for Level III ecoregions. There are fourteen aggregate nutrient ecoregions. Each aggregate nutrient ecoregion is divided into smaller ecoregions (subecoregions) referred to as Level III ecoregions. EPA will determine criteria for the waterbody types and Level III ecoregions within the following aggregate nutrient ecoregions:

- Lakes and Reservoirs
 - Aggregate Nutrient ecoregions: 3, 4, 5, and 14
- Rivers and Streams
 - Aggregate Nutrient ecoregions: 1, 4, 5, 8, and 10

1.2 References

This section lists documents that contain baselines, standards, guidelines, policies, and references that apply to the data analysis. Listed editions were valid at the time of publication. All documents are subject to revision, but these specific editions govern the concepts described in this document.

Nutrient Criteria Technical Guidance Document: Lakes and Reservoirs (Draft). EPA, Office of Water, EPA 822-D-99-001, April 1999.

Nutrient Criteria Technical Guidance Manual: Rivers and Streams (Draft). EPA, Office of Water, EPA 822-D-99-003, September 1999.

Guidance for Data Quality Assessment: Practical Methods for Data Analysis. EPA, Office of Research and Development, EPA QA/G-9, January 1998.

2.0 QA/QC PROCEDURES

In order to develop nutrient criteria, EPA needed to obtain nutrient data from the states. EPA requested nutrient data from the states and forwarded the data sets to INDUS via e-mail and/or US mail. In addition, EPA tasked INDUS to convert data from three national data sets. EPA

provided INDUS with a Legacy STORET extraction to convert into the database. The United States Geologic Survey (USGS) sent INDUS a CD-ROM with NASQAN data to convert. INDUS downloaded NAWQA files from the USGS Web site to convert the data. In total, INDUS converted and imported the following national and state data sets into the Nutrient Criteria Database:

- Legacy STORET
- NAWQA
- NASQAN
- EPA Region 1
- EPA Region 2 - Lake Champlain Monitoring Project
- EPA Region 2 - NYSDEC Finger Lakes Monitoring Program
- EPA Region 2 - NY Citizens Lake Assessment Program
- EPA Region 2 - Lake Classification and Inventory Survey
- EPA Region 2 - NYCDEP (1990-1998)
- EPA Region 2 - NYCDEP (Storm Event data)
- EPA Region 2 - New Jersey Nutrient Data (Tidal Waters)
- EPA Region 5
- EPA Region 3
- EPA Region 3 - Nitrite Data
- EPA Region 3 - Choptank River files
- EPA Region 4 - Tennessee Valley Authority
- EPA Region 7 - Central Plains Center for BioAssessment (CPCB)
- EPA Region 7 - REMAP
- EPA Region 2 - Delaware River Basin Commission (1990-1998)
- EPA Region 3 - PA Lake Data
- EPA Region 3 - University of Delaware
- EPA Region 10
- University of Auburn
- EPA Region 8 - MT and WY
- EPA Region 9
- Suffolk County
- NYCDEC
- NY Lakes Morphometry
- EPA Region 8 - South Dakota
- EPA Region 8 - Colorado Reservoir
- EPA Region 4
- EPA Region 10 - Lake Data
- EPA Region 7 - Central Plains Center for BioAssessment (CPCB) 2
- EPA Region 8 - North Dakota
- EPA Region 8 - Eagle River
- EPA Region 8 - Utah
- Florida

As part of the conversion process, INDUS performed a number of Quality Assurance/Quality Control (QA/QC) steps to ensure that the data were properly converted into the Nutrient Criteria Database. Sections 2.1 and 2.2 explain the steps performed by INDUS to convert the data.

2.1 National Data Sets

INDUS converted three national data sets into the Nutrient Criteria Database: Legacy STORET data, NASQAN data, and NAWQA data. A previous EPA contractor performed the extraction of Legacy STORET data and documented the QA/QC procedures used on the data. This documentation is included in Appendix A. INDUS performed minimal QA/QC on the Legacy STORET data set because the previous contractor completed the steps outlined in Appendix A. INDUS and EPA also agreed to convert the NAWQA and NASQAN data sets with minimal QA/QC on the assumption that the source agency, the USGS, QA/QC'd the data.

For each of the three national data sets, INDUS ran queries to determine if 1) samples existed without results and 2) if stations existed without samples. Per Task Order Project Officer (TOPO) direction, these records were deleted from the system. For analysis purposes, EPA determined that there was no need to keep station records with no samples and sample records with no results. INDUS also confirmed that each data set contained no duplicate records.

In addition, INDUS deleted all composite results from the Legacy STORET data. Per TOPO direction, it was decided that composite sample results would not be used in the statistical analysis.

2.2 State Data

Each state data set was delivered in a unique format. Many of the data sets were delivered to INDUS without corresponding documentation. INDUS analyzed each state data set in order to determine which parameters should be converted for analysis. INDUS obtained a master parameter table from EPA and converted the parameters in the state data sets according to those that were present in the EPA parameter table. INDUS converted all of the data elements in the state data sets that mapped directly to the Nutrient Criteria Database; data elements that did not map to the Nutrient Criteria Database were not converted. In some cases, state data elements that did not directly map into the Oracle database were inserted into a comment field within the database. Also, INDUS maintained an internal record of which state data elements were inserted into the comment field.

As part of the data clean-up efforts, INDUS determined whether or not there were any duplicate records in the state data sets and deleted the duplicate records. INDUS checked the waterbody, station, and sample entities for duplicate records. However, if there was not enough information provided to determine duplicates such as sampling date, there was no way for INDUS to locate duplicate records. In addition, INDUS deleted station records with no samples and sample records with no results. INDUS also deleted waterbody records that were not associated with a station. In each case, INDUS maintained an internal record of how many records were deleted.

If INDUS encountered referential integrity errors, such as samples that referred to stations that did not exist, or if INDUS was unsure of whether a record was a duplicate, INDUS contacted the agency directly via e-mail or phone to resolve any issues that arose. INDUS saved an electronic copy of each e-mail correspondence with the states to ensure that a record of the decision was maintained.

Finally, INDUS examined the remark codes of each result record in the state data sets. INDUS mapped the remark codes to the STORET remark codes listed in Table 2 of Appendix A. If any of the state result records were associated with remark codes marked as “Delete” in Table 2 of Appendix A, the result records were not converted into the database.

2.3 Laboratory Methods

Many of the state data sets did not contain laboratory method information. In addition, laboratory method information was not available for the three national data sets. In order to determine missing laboratory method information, EPA tasked another contractor to contact the data owners to obtain the laboratory method. In some cases, the data owners responded and the laboratory methods were added to the database. In other cases, the methods are unknown.

2.4 Waterbody Name and Class Information

A large percentage of the data did not have waterbody-specific information. The only waterbody information contained in the three national data sets was the waterbody name, which was embedded in the station ‘location description’ field. Most of the state data sets contained waterbody name information; however, much of the data were duplicated throughout the data sets. Therefore, the waterbody information was cleaned manually. For the three national data sets, the ‘location description’ field was extracted from the station table and moved to a temporary table. The ‘location description’ field was sorted alphabetically. Unique waterbodies were grouped together based on name similarity and whether or not the waterbodies fell within the same county, state, and waterbody type. Finally, the ‘location description’ field was edited to include only waterbody name information, not descriptive information. For example, 110 MILE CREEK AT POMONA DAM OUTFLOW, KS PO-2 was edited to 110 MILE CREEK. Also, if 100 MILE CREEK was listed ten times in New York, but in four different counties, four 100 MILE CREEK waterbody records were created.

Similar steps were taken to eliminate duplicate waterbody records in the state data sets. If a number of records had similar waterbody names and fell within the same state, county, and waterbody type, the records were grouped to create a unique waterbody record.

Most of the waterbody data did not contain depth, surface area, and volume measurements. EPA needed this information to classify waterbody types. EPA attempted to obtain waterbody class information from the states. EPA sent waterbody files to the regional coordinators and requested that certain class information be completed by each state. The state response was poor; therefore, EPA was not able to perform statistical analysis for the waterbody types by class.

2.5 Ecoregion Data

Aggregate nutrient ecoregions and Level III ecoregions were added to the database using the station latitude and longitude coordinates, the county centroid, or HUC (Hydrological Unit Code) centroid. If a station was lacking latitude and longitude coordinates and county information, the data were not included in the statistical analysis. Appendix B lists the steps taken to add the two ecoregion types (aggregate and Level III) to the Nutrient Criteria Database. The ecoregion names were pulled from aggregate nutrient ecoregion and Level III ecoregion Geographical Information System (GIS) coverages. In summary, the station latitude and longitude coordinates were used to determine the ecoregion under the following circumstances:

- The latitude and longitude coordinates fell within the county/state listed in the station table.
- The county data were missing.

The county centroid was used to determine the ecoregions under the following circumstances:

- The latitude and longitude coordinates were missing, but the state/county information was available.
- The latitude and longitude coordinates fell outside the county/state/HUC listed in the station table. The county information was assumed to be correct; therefore, the county centroid was used.

The HUC centroid was used to determine the ecoregions under the following circumstances:

- The latitude and longitude coordinates and county were missing, but the HUC information was available.

If the latitude and longitude coordinates fell outside the continental US county coverage file (i.e., the point fell in the ocean or Mexico/Canada), the nearest ecoregion was assigned to the station.

3.0 STATISTICAL ANALYSIS REPORTS

Aggregate nutrient ecoregion tables were created by extracting all observations for a specific aggregate nutrient ecoregion from the Nutrient Criteria Database. Then, the data were reduced to create tables containing only the yearly median values. To create these tables, the median value for each waterbody was calculated using all observations for each waterbody by Level III ecoregion, state, county, year, and season. Tables of decade median values were created from the yearly median tables by calculating the median for each waterbody by Level III ecoregion, state, county, decade and season.

The Data Source and the Remark Code reports were created using all observations (all reported values). All the other reports were created from either the yearly median tables or the decade median tables. In other words, the descriptive statistics and regressions were run using the median values for each waterbody and not the individual reported values.

Statistical analyses were performed under the assumption that this data set is a random sample. If this assumption cannot be verified, the observations may or may not be valid. Values below the 1st and 99th percentile were removed from the Legacy STORET database prior to the creation of the national database. Also, data were treated according to the Legacy STORET remark codes in Appendix A.

The following contains a list of each report and the purpose for creating each report:

- Data Source—Created to provide a count of the amount of data and to identify the source(s).
- Remark Codes—Created to provide a description of the data.
- Median of Each Waterbody by Year—This was an intermediate step performed to obtain a median value for each waterbody to be used in the yearly descriptive statistics reports and the regression models.
- Median of Each Waterbody by Decade—This was an intermediate step performed to obtain a median value for each waterbody to be used in the decade descriptive statistics.
- Descriptive Statistics—Created to provide EPA with the desired statistics for setting criteria levels.
- Regression Models—Created to examine the relationships between biological and nutrient variables.

Note: Separate reports were created for each season.

3.1 Data Source Reports

Data source reports were presented in the following formats:

- The number and percentage of data from each data source were summarized in tables for each aggregate nutrient ecoregion by season and waterbody type.
- The number and percentage of data from each data source were summarized in tables for each aggregate nutrient ecoregion for all seasons and waterbody type.
- The number and percentage of data from each data source were summarized in tables for each Level III ecoregion by season and waterbody type.

The 'Frequency' represents the number of data values from a specific data source for each parameter by data source. The 'Row Pct' represents the percentage of data from a specific data source for each parameter.

3.2 Remark Code Reports

Remark code reports were presented in the following formats:

- The number and percentage of data associated with a particular remark code for each parameter were summarized in tables by Level III ecoregion by decade and season.

- The number and percentage of data associated with a particular remark code for each parameter were summarized in tables by Level III ecoregion by year and season.

The 'Frequency' represents the number of data values corresponding to the remark code in the column. The 'Row Pct' represents the percentage of data that was associated with the remark code in that row.

In the database, remark codes that were entered by the states were mapped to Legacy STORET remark codes. Prior to the analysis, the data were treated according to these remark codes. For example, if the remark code was 'K,' then the reported value was divided by two. Appendix A contains a complete list of Legacy STORET remark codes.

Note: For the reports, a remark code of 'Z' indicates that no remark codes were recorded. It does not correspond to Legacy STORET code 'Z.'

3.3 Median of Each Waterbody

To reduce the data and to ensure heavily sampled waterbodies or years were not over represented in the analysis, median value tables (described above) were created. The yearly median tables and decade median tables were delivered to the EPA in electronic format as csv (comma separated value or comma delimited) files.

3.4 Descriptive Statistic Reports

The number of waterbodies, median, mean, minimum, maximum, 5th, 25th, 75th, 95th percentiles, standard deviation, standard error, and coefficient of variation were calculated. The tables (described above) containing the decade median values for each waterbody for each parameter were used to create descriptive statistics reports for:

- Level III ecoregions by decade and season
- Aggregate nutrient ecoregions by decade and season

In addition, the tables containing the yearly median values for each waterbody for each parameter were used to create descriptive statistics reports for:

- Level III ecoregions by year and season

3.5 Regression Models

Simple linear regressions using the least squares method were performed to examine the relationships between biological and nutrient variables in lakes and reservoirs, and rivers and streams. Regressions were performed using the yearly median tables. Chlorophyll(s) in micrograms per liter (ug/L), Secchi in meters (m), Dissolved Oxygen in milligrams per liter (mg/L), Turbidity, and pH were the biological variables in these models. Secchi data were used in the lake and reservoir models, and Turbidity data were used in the river and stream models.

The nutrient variables in these models include: Total Phosphorus in ug/L, Total Nitrogen in mg/L, Total Kjeldahl Nitrogen in mg/L, and Nitrate and Nitrite in mg/L.

4.0 TIME PERIOD

Data collected from January 1990 to December 2000 were used in the statistical analysis reports. To capture seasonal differences, the data were classified as follows:

- Aggregate nutrient ecoregions: 6, 7, and 8
 - Spring: April to May
 - Summer: June to August
 - Fall: September to October
 - Winter: November to March
- Aggregate nutrient ecoregions: 1, 2, 3, 4, 5, 9, 10, 11, 12, 13, and 14
 - Spring: March to May
 - Summer: June to August
 - Fall: September to November
 - Winter: December to February

5.0 DATA SOURCES AND PARAMETERS FOR THE AGGREGATE NUTRIENT ECOREGIONS

This section provides information for the nutrient aggregate ecoregions that were analyzed by waterbody type. Each section lists the data sources for the aggregate nutrient ecoregion including: 1) the data sources, 2) the parameters included in the analysis, and 3) the Level III ecoregions within the aggregate nutrient ecoregions.

Note: For analysis purposes, data for the following parameters were grouped together and reported under Phosphorous, Dissolved Inorganic (DIP):

Phosphorus, Dissolved Inorganic (DIP)
Phosphorus, Dissolved (DP)
Phosphorus, Dissolved Reactive (DRP)
Orthophosphate, dissolved, mg/L as P
Orthophosphate (OPO4_PO4)

5.1 Lakes and Reservoirs

5.1.1 Aggregate Nutrient Ecoregion 3

Data Sources:

Legacy STORET

EPA Region 10

EPA Region 8 - Colorado Reservoir

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)

Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)

Chlorophyll A, Trichromatic, uncorrected (ug/L)

Dissolved Inorganic Phosphorus (DIP) (ug/L)

Dissolved Oxygen (DO) (mg/L)

Nitrite and Nitrate, (NO₂+NO₃) (mg/L)

Nitrogen, Total (TN) (mg/L)

Nitrogen, Total Kjeldhal (TKN) (mg/L)

Phosphorus, Total (TP) (ug/L)

SECCHI (m)

pH

Level III ecoregions:

6, 10, 12, 13, 18, 20, 22, 24, 80, 81

5.1.2 Aggregate Nutrient Ecoregion 4

Data Sources:

Legacy STORET

EPA Region 8 - MT and WY

EPA Region 8 - South Dakota

EPA Region 8 - North Dakota

Parameters:

Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)

Chlorophyll A, Trichromatic, uncorrected (ug/L)

Dissolved Inorganic Phosphorus (DIP) (ug/L)

Dissolved Oxygen (DO) (% Saturated)

Dissolved Oxygen (DO) (mg/L)

Nitrite and Nitrate, (NO₂+NO₃) (mg/L)

Nitrogen, Total (TN) (mg/L)

Nitrogen, Total Kjeldhal (TKN) (mg/L)
Phosphorus, Total (TP) (ug/L)
SECCHI (m)
pH

Level III ecoregions:

26, 28, 30, 31, 43, 44

5.1.3 Aggregate Nutrient Ecoregion 5

Data sources:

Legacy STORET
EPA Region 8 - MT and WY
EPA Region 8 - South Dakota
EPA Region 8 - North Dakota

Parameters:

Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
Chlorophyll A, Trichromatic, uncorrected (ug/L)
Dissolved Inorganic Phosphorus (DIP) (ug/L)
Dissolved Oxygen (DO) (% Saturated)
Dissolved Oxygen (DO) (mg/L)
Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
Nitrogen, Total (TN) (mg/L)
Nitrogen, Total Kjeldhal (TKN) (mg/L)
Phosphorus, Total (TP) (ug/L)
SECCHI (m)
pH

Level III ecoregions:

25, 27, 32, 42

5.1.4 Aggregate Nutrient Ecoregion 14

Data sources:

Legacy STORET
Region 2 - NY Citizens Lake Assessment Program
Region 2 - NYCDEP (1990-1998)
EPA Region 1

Parameters:

CHLB (ug/L)
 CHLC (ug/L)
 Chlorophyll A, Fluorometric, corrected (ug/L)
 Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
 Chlorophyll A, Phytoplankton, spectrophotometric, uncorrected (ug/L)
 Chlorophyll A, Trichromatic, uncorrected (ug/L)
 Dissolved Inorganic Phosphorus (DIP) (ug/L)
 Dissolved Oxygen (DO) (mg/L)
 Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
 Nitrogen, Total (TN) (mg/L)
 Nitrogen, Total Kjeldhal (TKN) (mg/L)
 Phosphorus, Total (TP) (ug/L)
 SECCHI (m)
 pH

Level III ecoregions:

59, 63, 84

5.2 Rivers and Streams**5.2.1 Aggregate Nutrient Ecoregion 1**Data sources:

Legacy STORET
 NASQAN
 NAWQA
 EPA Region 10

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)
 Chlorophyll A, Periphyton, spectrophotometric, uncorrected (mg/sqm)
 Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
 Chlorophyll A, Trichromatic, uncorrected (ug/L)
 Dissolved Inorganic Phosphorus (DIP) (ug/L)
 Dissolved Oxygen (DO) (mg/L)
 Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
 Nitrogen, Total (TN) (mg/L)
 Nitrogen, Total Kjeldhal (TKN) (mg/L)
 Phosphorus, Total (TP) (ug/L)
 Phosphorus, orthophosphate, total, as P(ug/L)
 Turbidity (FTU)

Turbidity (NTU)
Turbidity (JCU)
pH

Level III ecoregions:

3, 7

5.2.2 Aggregate Nutrient Ecoregion 4

Data sources:

Legacy STORET
NASQAN
NAWQA
EPA Region 7 - Central Plains Center for BioAssessment (CPCB)
EPA Region 7 - Central Plains Center for BioAssessment (CPCB) 2
EPA Region 7 - REMAP
EPA Region 8 - MT and WY
EPA Region 8 - South Dakota
EPA Region 8 - North Dakota

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)
Chlorophyll A, Pheophytin, corrected (ug/L)
Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
Dissolved Inorganic Phosphorus (DIP) (ug/L)
Dissolved Oxygen (DO) (% Saturated)
Dissolved Oxygen (DO) (mg/L)
Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
Nitrogen, Total (TN) (mg/L)
Nitrogen, Total Kjeldhal (TKN) (mg/L)
Organic_P (ug/L)
Phosphorus, Total (TP) (ug/L)
Phosphorus, orthophosphate, total, as P(ug/L)
Turbidity (FTU)
Turbidity (NTU)
Turbidity (JCU)
pH

Level III ecoregions:

26, 28, 30, 31, 43, 44

5.2.3 Aggregate Nutrient Ecoregion 5

Data sources:

Legacy STORET

NASQAN

NAWQA

EPA Region 7 - Central Plains Center for BioAssessment (CPCB)

EPA Region 7 - Central Plains Center for BioAssessment (CPCB) 2

EPA Region 7 - REMAP

EPA Region 8 - MT and WY

EPA Region 8 - South Dakota

EPA Region 8 - North Dakota

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)

Chlorophyll A, Pheophytin, corrected (ug/L)

Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)

Dissolved Inorganic Phosphorus (DIP) (ug/L)

Dissolved Oxygen (DO) (% Saturated)

Dissolved Oxygen (DO) (mg/L)

Nitrite and Nitrate, (NO₂+NO₃) (mg/L)

Nitrogen, Total (TN) (mg/L)

Nitrogen, Total Kjeldhal (TKN) (mg/L)

Organic_P (ug/L)

Phosphorus, Total (TP) (ug/L)

Phosphorus, orthophosphate, total, as P (ug/L)

Turbidity (FTU)

Turbidity (NTU)

Turbidity (JCU)

pH

Level III ecoregions:

25, 27, 32, 42

5.2.4 Aggregate Nutrient Ecoregion 8

Data sources:

Legacy STORET

NASQAN

NAWQA

EPA Region 2 - NYCDEP (1990-1998)

EPA Region 1

EPA Region 3
EPA Region 5

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)
Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
Chlorophyll A, Phytoplankton, spectrophotometric, uncorrected (ug/L)
Chlorophyll A, Trichromatic, uncorrected (ug/L)
Dissolved Inorganic Phosphorus (DIP) (ug/L)
Dissolved Oxygen (DO) (% Saturated)
Dissolved Oxygen (DO) (mg/L)
Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
Nitrogen, Total (TN) (mg/L)
Nitrogen, Total Kjeldhal (TKN) (mg/L)
Phosphorus, Total (TP) (ug/L)
Phosphorus, orthophosphate, total, as P (ug/L)
Turbidity (FTU)
Turbidity (NTU)
pH

Level III ecoregions:

49, 50, 58, 62, 82

5.2.5 Aggregate Nutrient Ecoregion 10

Data sources:

Legacy STORET
NASQAN
EPA Region 7 - Central Plains Center for BioAssessment (CPCB)
EPA Region 7 - Central Plains Center for BioAssessment (CPCB) 2
EPA Region 7 - REMAP

Parameters:

Chlorophyll A, Fluorometric, corrected (ug/L)
Chlorophyll A, Pheophytin, corrected (ug/L)
Chlorophyll A, Phytoplankton, chromatographic- fluorometric (ug/L)
Chlorophyll A, Phytoplankton, spectrophotometric Acid (ug/L)
Chlorophyll A, Trichromatic, uncorrected (ug/L)
Chlorophyll B, Phytoplankton, chromatographic- fluorometric (ug/L)
Dissolved Inorganic Phosphorus (DIP) (ug/L)
Dissolved Oxygen (DO) (mg/L)
Nitrite and Nitrate, (NO₂+NO₃) (mg/L)
Nitrogen, Total (TN) (mg/L)

Nitrogen, Total Kjeldhal (TKN) (mg/L)
Organic_P (ug/L)
Phosphorus, Total (TP) (ug/L)
Phosphorus, orthophosphate, total, as P(ug/L)
Turbidity (FTU)
Turbidity (NTU)
Turbidity (JCU)
pH

Level III ecoregions:

34, 73

APPENDIX A. Process Used to QA/QC the Legacy STORET Nutrient Data Set

1. STORET water quality parameters and Station and Sample data items were retrieved from USEPA's mainframe computer. Table 1 lists all retrieved parameters and data items.

TABLE 1: PARAMETERS AND DATA ITEMS RETRIEVED FROM STORET		
Parameters Retrieved (STORET Parameter Code)	Station Data Items Included (STORET Item Name)	Sample Data Items Included (STORET Item Name)
TN - mg/l (600) TKN - mg/l (625) Total Ammonia (NH ₃ +NH ₄) - mg/l (610) Total NO ₂ +NO ₃ - mg/l (630) Total Nitrite - mg/l (615) Total Nitrate - mg/l (620) Organic N - mg/L (605) TP - mg/l (665) Chlor <i>a</i> - ug/L (spectrophotometric method, 32211) Chlor <i>a</i> - ug/L (fluorometric method corrected, 32209) Chlor <i>a</i> - ug/L (trichromatic method corrected, 32210) Secchi Transp. - inches (77) Secchi Transp. - meters (78) +Turbidity JCU's (70) +Turbidity FTU's (76) +Turbidity NTU's field (82078) +Turbidity NTU's lab (82079) +DO - mg/L (300) +Water Temperature (degrees C, 10/degrees F, 11)	Station Type (TYPE) Agency Code (AGENCY) Station No. (STATION) Latitude - std. decimal degrees (LATSTD) Longitude - std. decimal degrees (LONGSTD) Station Location (LOCNAME) County Name (CONAME) State Name (STNAME) Ecoregion Name - Level III (ECONAME) Ecoregion Code -Level III (ECOREG) Station Elevation (ELEV) Hydrologic Unit Code (CATUNIT) RF1 Segment and Mile (RCHMIL) RF1ON/OFF tag (ONOFF)	Sample Date (DATE) Sample Time (TIME) Sample Depth (DEPTH) Composite Sample Code (SAMPMETHOD)
+ If data record available at a station included data only for this or other such marked parameters, data record was deleted from data set.		

The following set of retrieval rules were applied to the retrieval process:

- Data were retrieved for waterbodies specified only as **'lake', 'stream', 'reservoir', or 'estuary'** under "Station Type" parameter. Any stations specified as 'well,' 'spring,' or 'outfall' were eliminated from the retrieved data set.
- Data were retrieved for station types described as 'ambient' (e.g., no pipe or facility discharge data) under the "Station Type" parameter.
- Data were retrieved that were designated as 'water' samples only. This includes 'bottom' and 'vertically integrated' water samples.
- Data were retrieved that were designated as either 'grab' samples and 'composite' samples (mean result only).

- No limits were specified for sample depths.
 - Data were retrieved for all fifty states, Puerto Rico, and the District of Columbia.
 - The time period specified for data retrieval was January 1990 to September 1998.
 - No data marked as “Retired Data” (i.e., data from a generally unknown source) were retrieved.
 - Data marked as “National Urban Runoff data” (i.e., data associated with sampling conducted after storm events to assess nonpoint source pollutants) were included in the retrieval. Such data are part of STORET’s ‘Archived’ data.
 - Intensive survey data (i.e., data collected as part of specific studies) were retrieved.
2. Any values falling below the 1st percentile and any values falling above the 99th percentile were transformed into ‘missing’ values (i.e., values were effectively removed from the data set, but were not permanently eliminated).
 3. Based on the STORET ‘Remark Code’ associated with each retrieved data point, the following rules were applied (Table 2):

TABLE 2: STORET REMARK CODE RULES	
STORET Remark Code	Keep or Delete Data Point
blank - Data not remarked.	Keep
A - Value reported is the mean of two or more determinations.	Keep
B - Results based upon colony counts outside the acceptable ranges.	Delete
C - Calculated. Value stored was not measured directly, but was calculated from other data available.	Keep
D - Field measurement.	Keep
E - Extra sample taken in compositing process.	Delete
F - In the case of species, F indicates female sex.	Delete
G - Value reported is the maximum of two or more determinations.	Delete
H - Value based on field kit determination; results may not be accurate.	Delete
I - The value reported is less than the practical quantification limit and greater than or equal to the method detection limit.	Keep, but used one-half the reported value as the new value.
J - Estimated. Value shown is not a result of analytical measurement.	Delete

TABLE 2: STORET REMARK CODE RULES	
K - Off-scale low. Actual value not known, but known to be less than value shown.	Keep, but used one-half the reported value as the new value.
L - Off-scale high. Actual value not known, but known to be greater than value shown.	Keep
M -Presence of material verified, but not quantified. Indicates a positive detection, at a level too low to permit accurate quantification.	Keep, but used one half the reported value as the new value.
N -Presumptive evidence of presence of material.	Delete
O -Sample for, but analysis lost. Accompanying value is not meaningful for analysis.	Delete
P -Too numerous to count.	Delete
Q -Sample held beyond normal holding time.	Delete
R -Significant rain in the past 48 hours.	Delete
S -Laboratory test.	Keep
T -Value reported is less than the criteria of detection.	Keep, but replaced reported value with 0.
U -Material was analyzed for, but not detected. Value stored is the limit of detection for the process in use.	Keep, but replaced reported value with 0.
V -Indicates the analyte was detected in both the sample and associated method blank.	Delete
W -Value observed is less than the lowest value reportable under remark "T."	Keep, but replaced reported value with 0.
X -Value is quasi vertically-integrated sample.	No data point with this remark code in data set.
Y -Laboratory analysis from unpreserved sample. Data may not be accurate.	Delete
Z -Too many colonies were present to count.	Delete
<p>If a parameter (excluding water temperature) value was less than or equal to zero and no remark code was present, the value was transformed into a missing value.</p> <p>Rationale - Parameter concentrations should never be zero without a proper explanation. A method detection limit should at least be listed</p>	

4. Station records were eliminated from the data set if any of the following descriptors were present within the “Station Type” parameter:
 - ▶ **MONITR** - Source monitoring site, which monitors a known problem or to detect a specific problem.
 - ▶ **HAZARD** - Site of hazardous or toxic wastes or substances.
 - ▶ **ANPOOL** - Anchialine pool, underground pools with subsurface connections to watertable and ocean.
 - ▶ **DOWN** - Downstream (i.e., within a potentially polluted area) from a facility which has a potential to pollute.
 - ▶ **IMPDMT** - Impoundment. Includes waste pits, treatment lagoons, and settling and evaporation ponds.
 - ▶ **STMSWR** - Storm water sewer.
 - ▶ **LNDFL** - Landfill.
 - ▶ **CMBMI** - Combined municipal and industrial facilities.
 - ▶ **CMBSRC** - Combined source (intake and outfall).

Rationale - these descriptors potentially indicate a station location that at which an ambient water sample would not be obtained (i.e., such sampling locations are potentially biased) or the sample location is not located within one of the designated water body types (i.e, ANPOOL).

5. Station records were eliminated from data set if the station location did not fall within any established cataloging unit boundaries based on their latitude and longitude.
6. Using nutrient ecoregion GIS coverage provided by USEPA, all station locations with latitude and longitude coordinates were tagged with a nutrient ecoregion identifier (nutrient region identifiers are values 1 - 14) and the associated nutrient ecoregion name. Because no nutrient ecoregions exist for Alaska, Hawaii, and Puerto Rico, stations located in these states were tagged with “dummy” nutrient ecoregion numbers (20 = Alaska, 21 = Hawaii, 22 = Puerto Rico).
7. Using information provided by TVA, 59 station locations that were marked as ‘stream’ locations under the “Station Type” parameter were changed to ‘reservoir’ locations.
8. The nutrient data retrieved from STORET were assessed for the presence of duplicate data records. The duplicate data identification process consisted of three steps: 1) identification of records that matched exactly in terms of each variable retrieved; 2) identification of records that matched exactly in terms of each variable retrieved except for their station identification numbers; and 3) identification of records that matched exactly in terms of each variable retrieved except for their collecting agency codes. The data duplication assessment procedures were conducted using SAS programs.

Prior to initiating the data duplication assessment process, the STORET nutrient data set contained:

41,210 station records
924,420 sample records

- Identification of exactly matching records

All data records were sorted to identify those records that matched exactly. For two records to match exactly, all variables retrieved had to be the same. For example, they had to have the same water quality parameters, parameter results and associated remark codes, and have the same station data item and sample data item information. Exactly matching records were considered to be exact duplicates, and one duplicate record of each identified matching set were eliminated from the nutrient data set. A total of 924 sample records identified as duplicates by this process were eliminated from the data set.

- Identification of matching records with the exception of station identification number

All data records were sorted to identify those records that matched exactly except for their station identification number (i.e., they had the same water quality parameters, parameter results and associated remark codes, and the same station and sample data item information with the exception of station identification number). Although the station identification numbers were different, the latitude and longitude for the stations were the same indicating a duplication of station data due to the existence of two station identification numbers for the same station. For each set of matching records, one of the station identification numbers was randomly selected and its associated data were eliminated from the data set. A total of 686 sample records were eliminated from the data set through this process.

- Identification of matching records with the exception of collecting agency codes

All data records were sorted to identify those records that matched exactly except for their collecting agency codes (i.e., they had the same water quality parameters, parameter results and associated remark codes, and the same station and sample data item information with the exception of agency code). The presence of two matching data records each with a different agency code attached to it suggested that one agency had utilized data collected by the other agency and had entered the data into STORET without realizing that it already had been placed in STORET by the other agency. No matching records with greater than two different agency codes were identified. For determining which record to delete from the data set, the following rules were developed:

- ▶ If one of the matching records had a USGS agency code, the USGS record was retained and the other record was deleted.
- ▶ Higher level agency monitoring program data were retained. For example, federal program data (indicated by a "1" at the beginning of the STORET agency code) were retained against state (indicated by a "2") and local (indicated by values higher than 2) program data.
- ▶ If two matching records had the same level agency code, the record from the agency with the greater number of overall observations (potentially indicating the data set as the source data set) was retained.

A total of 2,915 sample records were eliminated through this process.

As a result of the duplicate data identification process, a total of 4,525 sample records and 36 individual station records were removed from the STORET nutrient data set. The resulting

nutrient data set contains the following:

- 41,174 station records
- 919,895 sample records

APPENDIX B. Process for Adding Aggregate Nutrient Ecoregions and Level III Ecoregions

The flag_id tracks the type of changes that were made to the data. There are a total of eight flags that are used to describe the changes made to the data. The flags are defined as follows:

- 1—The latitude and longitude coordinates match the county that was provided. If the HUC was null, it was updated based on the latitude and longitude coordinates. The ecoregions were determined by using the latitude and longitude coordinates.
- 2—The county and HUC are available, but the latitude and/or longitude coordinates are missing. Therefore, the centroid of the intersection of the county and HUC was used to determine the ecoregions and the latitude and longitude coordinates. If the HUC and county did not intersect, the county centroid was used to determine the ecoregions and the latitude and longitude coordinates.
- 3—The county is available, but the HUC and the latitude and/or longitude coordinates are missing. Therefore, the county centroid was used to determine the ecoregions, HUC, and the latitude and longitude coordinates.
- 4—The HUC is available, but the county is not and the latitude and/or longitude coordinates are missing. Therefore, the HUC centroid was used to determine the ecoregions, county, and the latitude and longitude coordinates.
- 5—The county is missing, but the latitude and longitude coordinates are available. Note: A county is considered missing if it is invalid. In other words, if the county entered did not exist in the state, it was considered null. Therefore, the latitude and longitude coordinates were used to determine the ecoregions, county, and HUC (if it was missing).
- 6—The latitude and longitude coordinates did not match the county that was provided, but they did match the HUC. Therefore, the county centroid was used to determine ecoregion values.
- 7—The latitude and longitude coordinates did not match the county or the HUC that was provided (including null HUCs). Therefore, the county centroid was used to determine ecoregion values.
- 8—The latitude and longitude coordinates were missing, but the ecoregions were provided by the state. The ecoregions provided by the states were used as the ecoregion values.

APPENDIX C. Glossary

Coefficient of Variation - A measure of variability. The standard deviation divided by the mean multiplied by 100.

Maximum - The highest value.

Mean – A measure of central tendency. The arithmetic average.

Median – A measure of central tendency. The value which cuts the distribution in half, such that half of the values are above the median, and half of the values are below the median. Also called the 50th percentile or middle value.

Minimum - The lowest value.

Standard Deviation – A measure of variability. The square root of the variance with the variance defined as the sum of the squared deviations divided by the sample size minus one.

Standard Error - A measure of variability. The standard deviation divided by the square root of the sample size.

5th % - the 5th percentile

25th % - the 25th percentile, the first quartile.

75th % - the 75th percentile, the third quartile.

95th % - the 95th percentile